

## Species List of the Diptera of the Outer

### Hebrides (Revised January 2020)

#### Introduction

The first comprehensive review of the Diptera of the Outer Hebrides was included in *Diptera Scotica VI – The Western Isles* (Grimshaw 1914-1916) and produced a list of 251 species for the Outer Hebrides. The attraction of the Outer Hebrides, and St Kilda in particular, to naturalists did not diminish and some 65 years later, A.R. Waterston was able to compile a list of 591 species (Waterston 1981).

The next annotated species checklist also combined the islands of the Outer Hebrides and the Ebudes (Inner Hebrides) and was based on field work by Peter Skidmore from 1960-1995, literature searches and specimens in museum and private collections. Overall 1,895 species are listed, including 769 species from the Outer Hebrides (Skidmore 2008).

The list compiled by Skidmore is the base-line for this revised list, supplemented by species listed on the NBN Atlas and records in the scientific literature. The taxonomy has been up-dated so that it is consistent with the latest version of the UK species list (Chandler 2019). The list now includes 841 species (11.7% of the British species) from 64 of the 107 families present in the UK.

The records incorporated in the list are based on survey data from a large number of sources, comprising both casual records and systematic surveys, collected from a variety of habitats using a range of sampling methods and at different times of year. The diversity of species recorded is also influenced by the ease of recording and identification, and the interests of specialist recorders working in the area. For example, large and morphologically distinctive species such as hoverflies are more commonly recorded than fungus gnats, which are small dark flies usually less than 2mm long. Therefore, considerable care is required when interpreting the information derived from the checklist as it contains temporal, geographical and sampling biases. This is a common feature of many taxonomic checklists where a large proportion of the data are derived from a wide range of sources.

Waterston (1981) and Skidmore do not quote their sources for each species and provide little additional information other than the islands where a species has been recorded. Therefore, for some species it has not been possible to include the original source. Additional information will be added as the research progresses.

Skidmore (2008) includes some unpublished material and these records are included as “cited by”. Some of the published material included sufficient information to form a coherent biological record and these are included in the OHBR dataset “Outer Hebrides Insect Records from Literature” on the [National Biodiversity Network Atlas](#).

The sources also include references to records on the NBN Atlas from other organisations e.g. Biological Records Centre (BRC). It should be noted that there are differences in the nomenclature used on the NBN Atlas and the current edition of the UK Diptera Checklist (Chandler 2019). The use of a different synonym on NBN Atlas is recorded in the comments for each species.

Checklists are always a work in progress, and subject to revision as new species are recorded, and “doubtful” species either confirmed or removed from the list. Comments on the status of any species on the list or additional information is always appreciated.

A [bibliography](#) of sources accompanies this list and a more extensive list of literature relating to the natural history of the Outer Hebrides can be found on the [Outer Hebrides Natural History Bibliography](#) website.

Christine Johnson  
Outer Hebrides Biological Recording

## Tipuloidea: Tipulidae - Long-palped Craneflies

<i>Dolichopeza albipes</i>	(Ström, 1768)	St Kilda records only	Skidmore (2008); NBN: BRC 1974 (?)
<i>Nephrotoma appendiculata</i>	(Pierre, 1919)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013-2018
<i>Nephrotoma cornicina</i>	(Linnaeus, 1758)		Skidmore (1994), Skidmore (2008); NBN: BRC 1970, OHBR 2019
<i>Nephrotoma flavescens</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936) as <i>Pachyrrhina lineata</i> , Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Nephrotoma submaculosa</i>	Edwards, 1928		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR BRC 1970-2002, 2012-2017
<i>Prionocera turcica</i>	(Fabricius, 1787)		Whitely, Garland, Hancock (1994), Skidmore (2008): NBN: BRC 1969, 1988
<i>Tipula alpium</i>	Bergroth, 1888		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: BRC 1974-1981, OHBR 2018, BRC 1974-1981
<i>Tipula cheethami</i>	Edwards, 1924		Skidmore (2008)
<i>Tipula confusa</i>	van der Wulp, 1883		Evans (1915) as <i>T. marmorata</i> Butt Lewis lighthouse, Waterston (1981) as <i>T. marmorata</i> , Skidmore (1994), Skidmore (2008) cites Bratton 1987; NBN: BRC 1995-1997, OHBR 2019
<i>Tipula fulvipennis</i>	De Geer, 1776		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Tipula lateralis</i>	Meigen, 1804		Forrest, Waterston & Watson (1936), Skidmore (1994) Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013, 2019: BRC 1970-1988
<i>Tipula limbata</i>	Zetterstedt, [1838]	OHBR 2018 first record	NBN: OHBR 2018
<i>Tipula lunata</i>	Linnaeus, 1758		Skidmore (2008)
<i>Tipula maxima</i>	Poda, 1761	OHBR 2018 first record	NBN: OHBR 2018
<i>Tipula melanoceros</i>	Schummel, 1833	OHBR 2018 first record	NBN: OHBR 2018
<i>Tipula montana</i>	Curtis, 1834	OHBR 2013 first record	NBN: OHBR 2013
<i>Tipula oleracea</i>	Linnaeus, 1758		Lack (1932), Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: BRC 1962-1989, OHBR 2013-2017
<i>Tipula pagana</i>	Meigen, 1818	OHBR 2012 first record	NBN: OHBR 2012-2019
<i>Tipula paludosa</i>	Meigen, 1830		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997; NBN: BRC 1962-1997, OHBR 2013-2019
<i>Tipula rufina</i>	Meigen, 1818		Evans (1915) Butt Lewis lighthouse, Skidmore (1994), Hamilton (1963), Skidmore (2008); NBN: BRC 1974-1979, OHBR 2010-2019
<i>Tipula scripta</i>	Meigen, 1830		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: BRC 1974, 1988
<i>Tipula subcunctans</i>	Alexander, 1921		Waterston (1981) as <i>Tipula czizeki</i> , (Skidmore 2008); NBN: BRC 1998, 2002(?)
<i>Tipula subnodicornis</i>	Zetterstedt, [1838]		Skidmore (2008); NBN: BRC 1975

Tipuloidea: Tipulidae - Long-palped Craneflies			
<i>Tipula variicornis</i>	Schummel, 1833		Skidmore (2008) cites Roper 1995; NBN: BRC 1975
<i>Tipula varipennis</i>	Meigen, 1818		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: BRC 1974-1998
Tipuloidea: Pediciidae - Hairy-eyed Craneflies			
<i>Dicranota exclusa</i>	(Walker, 1848)		Skidmore (2008)
<i>Pedicia rivosa</i>	(Linnaeus, 1758)		Skidmore (2008); NBN: OHBR 2012-2019
<i>Tricyphona immaculata</i>	(Meigen, 1804)		Skidmore (2008) cites Barron 1987, Skidmore 1994; NBN: BRC 1962, 1997, OHBR 2013, 2018-2019
Tipuloidea: Limoniidae - Short-palped Craneflies			
<i>Antocha vitripennis</i>	(Meigen, 1830)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Austrolimnophila ochracea</i>	(Meigen, 1804)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Dicranomyia autumnalis</i>	(Staeger, 1840)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Bratton 1987, Roper 1995; NBN: BRC 1970 - Monachs, OHBR 2010-2013, BRC 1970 - Monachs
<i>Dicranomyia chorea</i>	(Meigen, 1818)		Waterston (1981) as <i>Limonia chorea</i> , Skidmore (2018); NBN: BRC 2002, 1973; OHBR 2013 Mingulay
<i>Dicranomyia didyma</i>	(Meigen, 1804)	OHBR 2019 first record	NBN: OHBR 2019
<i>Dicranomyia goritiensis</i>	(Mik, 1864)	OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<i>Dicranomyia mitis</i>	(Meigen, 1830)		Skidmore (2008) cites Roper 1995; NBN: OHBR 2013 - Mingulay
<i>Dicranomyia modesta</i>	(Meigen, 1818)		Waterston (1981) as <i>Limonia modesta</i> , Skidmore (1994), Skidmore (2008)
<i>Dicranomyia morio</i>	(Fabricius, 1787)		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008); NBN: BRC 1969 - Monachs
<i>Dicranomyia sera</i>	(Walker, 1848)	St Kilda records only	Skidmore (2008)
<i>Dicranomyia stigmatica</i>	(Meigen, 1830)		Skidmore (2008)
<i>Dicranomyia ventralis</i>	(Schummel, 1829)		Skidmore (2008); NBN: BRC 1969 - Monachs, OHBR 2019
<i>Dicranophragma nemorale</i>	(Meigen, 1818)		Whitely, Garland, Hancock (1994) as <i>Limnophila nemorale</i> , Skidmore (2008); NBN: BRC 1996, OHBR 2013- Mingulay, 2019
<i>Dicranophragma separatum</i>	(Walker, 1848)		Whitely, Garland, Hancock (1994) as <i>Limnophila separatum</i> , Skidmore (1994), Skidmore (2008)
<i>Erioconopa diuturna</i>	(Walker, 1848)		Skidmore (2008)
<i>Erioconopa trivialis</i>	(Meigen, 1818)		Lack (1932) St Kilda, Whitely, Garland, Hancock (1994) as <i>Ericoptera trivialis</i> , Skidmore (1994), Skidmore (2008)
<i>Erioptera fuscipennis</i>	Meigen, 1818		Lack (1932) St Kilda, Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: BRC 1970- Monachs

Tipuloidea: Limoniidae - Short-palped Craneflies			
<i>Erioptera lutea</i>	Meigen, 1804		Skidmore (2008) cites Roper 1995, BRC 1952 Broomfield & Vardy, OHBR 2019
<i>Euphylidorea meigenii</i>	(Verrall, 1886)		Lack 1931 St Kilda, Whitely, Garland, Hancock (1994) as <i>Limnophila meigenii</i> , Skidmore (2008); NBN: BRC 1962 -2002, OHBR 2010
<i>Euphylidorea phaeostigma</i>	(Schummel, 1829)	St Kilda records only	Skidmore (2008)
<i>Gonomyia conoviensis</i>	Barnes, 1924	OHBR 2013 first record	NBN: OHBR 2013
<i>Gonomyia dentata</i>	de Meijere, 1920		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Limonia nubeculosa</i>	Meigen, 1804		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: BRC 2002, OHBR 2018
<i>Limonia trivittata</i>	(Schummel, 1829)		Forrest, Waterston & Watson (1936) as <i>Limnobia trivittata</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Molophilus ater</i>	(Meigen, 1804)	OHBR 2018 first record	NBN: OHBR 2018
<i>Molophilus griseus</i>	(Meigen, 1804)		Skidmore (2008)
<i>Molophilus obscurus</i>	(Meigen, 1818)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: BRC 1996, OHBR 2013 Mingulay
<i>Molophilus occultus</i>	de Meijere, 1918		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Molophilus pleuralis</i>	de Meijere, 1920		Skidmore (2008)
<i>Molophilus propinquus</i>	(Egger, 1863)		Skidmore (2008)
<i>Ormosia hederæ</i>	(Curtis, 1831)		Lack (1931), Skidmore (2008)
<i>Ormosia nodulosa</i>	(Macquart, 1826)		Skidmore (2008)
<i>Ormosia pseudosimilis</i>	(Lundström, 1912)		Whitely, Garland, Hancock (1994), Skidmore 2008
<i>Ormosia staegeriana</i>	Alexander, 1953		Skidmore (2008)
<i>Phylidorea ferruginea</i>	(Meigen, 1818)		Skidmore (1995), Skidmore (2008); NBN: BRC 1970, 1989, OHBR 2013 - Mingulay
<i>Phylidorea fulvonervosa</i>	(Schummel, 1829)		Whitely, Garland, Hancock (1994) as <i>Limnophila fulvonervosa</i> , Skidmore (2008) cites Roper 1995; NBN: BRC 1962 (Broomfield & Vardy)
<i>Phylidorea squalens</i>	(Zetterstedt, [1838])	BRC 1996 first Record	NBN: BRC 1996
<i>Symplecta stictica</i>	(Meigen, 1818)		Whitely, Garland, Hancock (1994) as <i>Erioptera stictica</i> , Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Tasiocera murina</i>	(Meigen, 1818)		Skidmore (2008) cites Roper 1995, Bratton 1987
Bibionoidea: Bibionidae - St Mark's Flies			
<i>Biblio clavipes</i>	Meigen, 1818	<b>Requires review</b>	Skidmore (2008) - doubtful. Grimshaw (1914) record from Dale (1883). Old specimen needs to be checked. Included by Skartveit, Whittington & Bland (2013)

<b>Bibionoidea: Bibionidae - St Mark's Flies</b>			
<i>Bibio johannis</i>	(Linnaeus, 1767)	OHBR 2014 first record	NBN: OHBR 2014
<i>Bibio marci</i>	(Linnaeus, 1758)	OHBR 2016 first record	NBN: OHBR 2016, 2019
<i>Bibio nigriventris</i>	Haliday, 1833		Skartveit, Whittington & Bland (2013) - Broomfield & Vardy 1962 Lewis
<i>Bibio pomonae</i>	(Fabricius, 1775)		Waterston (1936), Grimshaw (1914) cites Dale 1882, Skidmore (2008), Skartveit, Whittington & Bland (2013); NBN: OHBR 2012-2018
<i>Dilophus febrilis</i>	(Linnaeus, 1758)		Skidmore (1994), Skidmore (2008), Skartveit, Whittington & Bland (2013) cite Broomfield & Vardy 1962; NBN: OHBR 2012-2017
<i>Dilophus femoratus</i>	Meigen, 1804		MacGraw (1994), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995, Skartveit, Whittington & Bland (2013); NBN: OHBR 2010-2015
<b>Sciarioidea: Keroplatidae - Fungus Gnats</b>			
<i>Macrocera centralis</i>	Meigen, 1818		Skidmore (2008)
<i>Macrocera fasciata</i>	Meigen, 1804		Skidmore (1994), Skidmore (2008)
<i>Macrocera stigma</i>	Curtis, 1837		Skidmore (1994), Skidmore (2008)
<b>Sciarioidea: Mycetophilidae - Fungus Gnats</b>			
<i>Acnemia nitidicollis</i>	(Meigen, 1818)	OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<i>Allodia lugens</i>	(Wiedemann, 1817)		Skidmore (1994), Skidmore (2008)
<i>Boletina dubia</i>	(Meigen, 1804)		Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Boletina plana</i>	Walker, 1856	St Kilda records only	Skidmore (2008)
<i>Brevicornu nigrofusum</i>	(Lundström, 1909)	OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<i>Brevicornu sericoma</i>	(Meigen, 1830)		Skidmore (1994), Skidmore (2008)
<i>Exechia fusca</i>	(Meigen, 1804)		Skidmore (1994), Skidmore (2008)
<i>Leia fascipennis</i>	Meigen, 1818		Skidmore (1994), Skidmore (2008)
<i>Mycetophila unipunctata</i>	Meigen, 1818		Skidmore (2008)
<i>Mycetophila britannica</i>	Laštovka & Kidd, 1975	Barra records only	Waterston (1981) as <i>M. lineola</i> , Skidmore (2008)
<i>Mycomya fimbriata</i>	(Meigen, 1818)	St Kilda records only	Skidmore (2008)
<i>Rymosia bifida</i>	(Edwards, 1925)		Skidmore (1994), Skidmore (2008)
<i>Rymosia fasciata</i>	(Meigen, 1804)		Skidmore (1994), Skidmore (2008)
<i>Sciophila hirta</i>	Meigen, 1818	St Kilda records only	Skidmore (2008)

Sciarioidea: Mycetophilidae - Fungus Gnats				
<i>Synapha fasciata</i>	Meigen, 1818	St Kilda records only	Skidmore (2008) cites a 1961 record from Hirta	
<i>Zygomyia humeralis</i>	(Wiedemann, 1817)		Skidmore (1994), Skidmore (2008)	
Sciarioidea: Sciaridae - Black fungus-gnats				
<i>Austrosciara hyalipennis</i>	(Meigen, 1804)		Waterston (198) as <i>Sciara autumnali</i> , Skidmore (2008) as <i>Ctenosciara hyalipennis</i> cites Buckland 1985 archaeological excavation	
<i>Bradysia nitidicollis</i>	(Meigen, 1818)		Waterston (198) as <i>Sciara nitidicollis</i> , Skidmore (2008)	
<i>Bradysia normalis</i>	Frey, 1948		Skidmore (2008)	
<i>Corynoptera postglobiformis</i>	Mohrig, 1993		Skidmore (2008)	
<i>Cratyna nobilis</i>	(Winnertz, 1867)		Skidmore (2008)	
<i>Phytosciara flavipes</i>	(Meigen, 1804)		Waterston (198) as <i>Sciara flavipes</i> , Skidmore (2008)	
<i>Trichosia morio</i>	(Fabricius, 1794)		Waterston (198) as <i>Trichosia caudata</i> , Skidmore (2008)	
Sciarioidea: Cecidomyiidae - Gall-midges				
<i>Anisostephus betulinus</i>	(Kieffer, 1889)		Skidmore (2008)	
<i>Asphondylia ulicis</i>	Trail, 1873		Skidmore (2008)	
<i>Bayeriola thymicola</i>	(Kieffer, 1888)	OHBR 2013 first record	NBN: OHBR 2013, 2014, 2016	
<i>Campylomyza flavipes</i>	Meigen, 1818		Skidmore (2008)	
<i>Contarinia barbichei</i>	(Kieffer, 1890)		Skidmore (2008) noted that <i>the presence of this species required review</i> (Heslop Harrison 1948). Species confirmed in 2014 (OHBR)	
<i>Contarinia jacobaeae</i>	(Loew, 1850)		Skidmore (2008)	
<i>Contarinia petioli</i>	(Kieffer, 1898)		Skidmore (2008); NBN: OHBR 2013, 2014	
<i>Contarinia tremulae</i>	Kieffer, 1909		Skidmore (2008)	
<i>Dasineura affinis</i>	(Kieffer, 1886)		Skidmore (2008); NBN: OHBR 2017	
<i>Dasineura angelicae</i>	Rübsaamen, 1916		Skidmore (2008)	
<i>Dasineura aucupariae</i>	(Kieffer, 1909)		Skidmore (2008)	
<i>Dasineura dioicae</i>	(Rübsaamen, 1895)		Skidmore (2008)	
<i>Dasineura engstfeldi</i>	(Rübsaamen, 1889)		Skidmore (2008)	
<i>Dasineura galiicola</i>	(Kieffer, 1909)		Skidmore (2008); NBN: OHBR 2017, 2018	
<i>Dasineura hygrophila</i>	(Mik, 1883)		Skidmore (2008)	
<i>Dasineura kiefferiana</i>	Marchal, 1896		Skidmore (2008)	
<i>Dasineura marginemtorquens</i>	(Brems, 1847)	<b>Requires confirmation</b>	Waterston (1981) as <i>Rabdophaga marginemtorquen</i> , Skidmore (2008) - doubtful - see Heslop Harrison (1948)	

Sciarioidea: Cecidomyiidae - Gall-midges			
<i>Dasineura plicatrix</i>	(Kieffer, 1909)		Skidmore (2008)
<i>Dasineura populeti</i>	(Rübsaamen, 1889)		Skidmore (2008)
<i>Dasineura pteridicola</i>	(Kieffer, 1901)		Skidmore (2008)
<i>Dasineura pustulans</i>	(Rübsaamen, 1889)		Skidmore (2008), Heslop Harrison (1948); NBN: OHBR 2014-2016
<i>Dasineura rosae</i>	(Bremi, 1847)		Skidmore (2008); NBN: OHBR 2018
<i>Dasineura ulmaria</i>	(Bremi, 1847)		Skidmore (2008); NBN: OHBR 2013, 2015
<i>Dasineura urticae</i>	(Perris, 1840)		Heslop Harrison (1948), Skidmore (2008); NBN: OHBR 2013-2018
<i>Geocrypta galii</i>	(Loew, 1850)	OHBR 2013 first record	NBN: OHBR 2013, 2015
<i>Harmandiola globuli</i>	(Rübsaamen, 1889)	OHBR 2014 first record	NBN: OHBR 2014
<i>Iteomyia capreae</i>	(Winnertz, 1853)		Heslop Harrison (1948), Skidmore (2008); NBN: OHBR 2014-2018
<i>Iteomyia major</i>	(Kieffer, 1889)		Skidmore (2008); NBN: OHBR 2013-2018
<i>Janetiella thymi</i>	(Kieffer, 1888)		Heslop Harrison (1940), Waterston (1981) and Skidmore (2008) possibly cites Heslop Harrison's record from Fiaray (Sound of Barra)
<i>Jaapiella alpina</i>	(F. Löw, 1885)	OHBR 2019 first record	NBN: OHBR 2019
<i>Massalongia rubra</i>	(Kieffer, 1890)		Skidmore (2008)
<i>Oligotrophus panteli</i>	Kieffer, 1898	HBRG 1997 first record	NBN: HBRG 1997 - Galls on <i>Juniperus communis alpina</i> . HBRG Juniper Gall Survey, 1995-2006 PF Entwistle
<i>Oligotrophus valerii</i>	(Tavares, 1904)		Skidmore (2008)
<i>Rhopalomyia ptarmicae</i>	(Vallot, 1849)		Skidmore (2008)
<i>Rabdophaga albipennis</i>	(Loew, 1850)	<b>Requires confirmation</b>	Heslop Harrison (1941) Heslop Harrison (1948), Skidmore (2008) - Records based on material from <i>Salix aurita</i> and <i>S. repens</i> collected by Heslop Harrison, but this species found only in terminal buds of <i>S. alba</i> .
<i>Rabdophaga cinerearum</i>	(Hardy, 1850)	OHBR 2017 first record	NBN: OHBR 2017. Synonymised with <i>R. strobilina</i> (Chandler 2019) but the species status is retained on NBN
<i>Rabdophaga dubiosa</i>	Kieffer, 1913	OHBR 2017 first record	NBN: OHBR 2017
<i>Rabdophaga karschi</i>	(Kieffer, 1891)	OHBR 2018 first record	NBN: OHBR 2018
<i>Rabdophaga pulvini</i>	(Kieffer, 1891)	OHBR 2018 first record	NBN: OHBR 2018
<i>Rabdophaga rosaria</i>	(Loew, 1850)	OHBR 2014 first record	NBN: OHBR 2014, 2017. Synonymised with <i>R. strobilina</i> on NBN Atlas, but now recognised as a separate species (Chandler 2019)
<i>Rabdophaga strobilina</i>	(Bremi, 1847)		Heslop Harrison (1948), Skidmore (2008); NBN: OHBR 2014, 2017-2018



Sciarioidea: Cecidomyiidae - Gall-midges			
<i>Semudobia tarda</i>	Roskam, 1977	OHBR 2014 first record	NBN: OHBR 2014 first record
Psychodoidea: Psychodidae - Owlet-midges, Mothflies			
<i>Pericoma pseudoexquisita</i>	Tonnoir, 1940		Waterston (1981) as <i>P. avicularia</i> , Skidmore (2008)
<i>Psychoda albipennis</i>	Zetterstedt, 1850	Only recorded on the Flannan Islands	Skidmore (2008)
<i>Psychoda trinouloosa</i>	Tonnoir, 1922		Skidmore (2008)
Trichoceroidea: Trichoceridae - Window-gnats			
<i>Trichocera regelationis</i>	(Linnaeus, 1758)		Skidmore (1994), Skidmore (2008) cites Roper 1995
Anisopodoidea: Anisopodidae - Window-gnats			
<i>Sylvicola cinctus</i>	(Fabricius, 1787)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997, Roper 1995; NBN: OHBR 2014, 2018
<i>Sylvicola fenestralis</i>	(Scopoli, 1763)	<b>Requires review</b>	Forrest, Waterson & Watson (1936) as <i>Phyrnus (Rhyphus) fenestralis</i> , Skidmore (2008) includes this species as doubtful
<i>Sylvicola punctatus</i>	(Fabricius, 1787)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2018
<i>Sylvicola zetterstedti</i>	(Edwards, 1923)		Whitely, Garland, Hancock (1994), Skidmore (2008)
Scatopsoidea: Scatopsidae - Black Scavenger Flies			
<i>Scatopse notata</i>	(Linnaeus, 1758)		Skidmore (2008); NBN: OHBR 2013
<i>Thripomorpha halterata</i>	(Meigen, 1838)	<b>Requires review</b>	Waterston (1981) as <i>Rhegmoclema halterat</i> , Skidmore (2008) refers to Freeman & Lane (1985) - only from Scottish Highlands
<i>Thripomorpha verralli</i>	(Edwards, 1934)	<b>Requires review</b>	Skidmore (2008) refers to Freeman & Lane (1985) - southern species. NBN lists 1984 record from Skye (HBRG MacGowan)
Ptychopteroidea: Ptychopteridae - Phantom Craneflies			
<i>Ptychoptera albimana</i>	(Fabricius, 1787)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2014-2016, 2019
<i>Ptychoptera scutellaris</i>	Meigen, 1818		Skidmore (2008)
Culicoidea: Dixidae - Meniscus-midges			
<i>Dixella aestivalis</i>	(Meigen, 1818)		Skidmore (2008)
<i>Dixella martinii</i>	(Peus, 1934)		Vardy & Broomfield 1962 - Lewis
Chironomoidea: Thaumaleidae - Trickle-midges			
<i>Thaumalea testacea</i>	Ruthe, 1831		Skidmore (2008) cites Beaumont 1994,



Chironomoidea: Thaumaleidae - Trickle-midges			
<i>Thaumalea verralli</i>	Edwards, 1929		Edwards & Collin (1932), Skidmore (2008) cites Beaumont 1994
Chironomoidea: Simuliidae - Blackflies			
<i>Simulium aureum</i>	Fries, 1824		Skidmore (2008)
<i>Simulium intermedium</i>	Roubaud, 1906		Waterston & Lyster (1979), Waterston (1981) as <i>S. nitidifrons</i> , Skidmore (2008)
<i>Simulium latipes</i>	(Meigen, 1804)	<b>Requires review</b>	Waterston (1981), not listed by Skidmore (2008)
<i>Simulium ornatum</i>	Meigen, 1818		Skidmore (2008)
<i>Simulium reptans</i>	(Linnaeus, 1758)		Skidmore (2008)
<i>Simulium vernalis</i>	Macquart, 1826		Skidmore (2008)
Chironomoidea: Ceratopogonidae - Biting-midges			
<i>Atrichopogon pavidus</i>	(Winnertz, 1852)		Skidmore (2008)
<i>Brachypogon sociabilis</i>	(Goetghebuer, 1920)		Waterston (1981) as <i>Isohelea sociabilis</i> , Skidmore (2008)
<i>Culicoides impunctatus</i>	Goetghebuer, 1920	<b>Requires review</b>	Skidmore (2008); NBN: iRecord 2019 (unconfirmed)
<i>Culicoides kibunensis</i>	Tokunaga, 1937		Skidmore (2008)
<i>Culicoides obsoletus</i>	(Meigen, 1818)	St Kilda record(s) only	Skidmore (2008)
<i>Culicoides pictipennis</i>	(Staeger, 1839)	St Kilda record(s) only	Skidmore (2008)
<i>Culicoides pulicaris</i>	(Linnaeus, 1758)		Skidmore (2008)
<i>Dasyhelea modesta</i>	(Winnertz, 1852)		Skidmore (2008) & Waterston (1981) as <i>D. holosericea</i> and <i>D. aestiva</i> synonyms
<i>Forcipomyia titillans</i>	(Winnertz, 1852)		Skidmore (2008)
<i>Palpomyia flavipes</i>	(Meigen, 1804)		Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Palpomyia nigripes</i>	(Meigen, 1830)		Skidmore (2008)
<i>Palpomyia quadispinosa</i>	Goetghebuer, 1920		Whitely, Garland, Hancock (1994), Skidmore (2008): NBN: iRecord - Whitely 1989, presumably the same record?
<i>Palpomyia spinipes</i>	(Meigen in Panzer, 1806)		Skidmore (2008)
<i>Schizohela leucopeza</i>	(Meigen, 1804)		Waterston (1981) as <i>Monohelea leucopeza</i> , Skidmore (2008)
<i>Serromyia femorata</i>	(Meigen, 1804)		Skidmore (2008)
<i>Serromyia morio</i>	(Fabricius, 1775)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: iRecord -Whitely 1989 - presumably the same record from Eriskay?
Chironomoidea: Chironomidae - Non-biting Midges			
<i>Ablabesmyia monilis</i>	(Linnaeus, 1758)		Skidmore (2008)

### Chironomoidea: Chironomidae - Non-biting Midges

<i>Acricotopus lucens</i>	(Zetterstedt, 1850)		Skidmore (2008)
<i>Arctopelopia griseipennis</i>	(van der Wulp, 1859)		Skidmore (2008)
<i>Brillia bifida</i>	(Kieffer, 1909)		Waterston (1981) as <i>B. modesta</i> , Skidmore (2008)
<i>Bryophaenocladus vernalis</i>	(Goetghebuer, 1921)		Skidmore (2008)
<i>Chaetocladus dissipatus</i>	(Edwards, 1929)	Flannan Islands only	Skidmore (2008) and Waterston (1981) Flannan Islands
<i>Chaetocladus melaleucus</i>	(Meigen, 1818)	St Kilda records only	Skidmore (2008)
<i>Chaetocladus perennis</i>	(Meigen, 1830)		Skidmore (2008)
<i>Chironomus aprilius</i>	Meigen, 1830		Skidmore (2008)
<i>Chironomus longistylus</i>	Goetghebuer, 1921		Skidmore (2008)
<i>Chironomus plumosus</i>	(Linnaeus, 1758)		Waterston & Lyster (1979), Skidmore (2008). Also listed as <i>C. annularis</i> synonym by Skidmore (2008) and Waterston (1981)
<i>Chironomus salinarius</i>	Kieffer in Thienemann, 1915		Skidmore (2008) cites Buckland 1988 (archaeological dig)
<i>Chironomus tentans</i>	Fabricius, 1805		Skidmore (2008)
<i>Cladopelma viridulum</i>	(Linnaeus, 1767)		Waterston & Lyster (1979), Skidmore (2008),
<i>Clunio marinus</i>	Haliday, 1855		Skidmore (2008)
<i>Conchapelopia melanops</i>	(Meigen, 1818)		Skidmore (2008)
<i>Corynoneura scutellata</i>	Winnertz, 1846		Skidmore (2008)
<i>Cricotopus annulator</i>	Goetghebuer, 1927		Skidmore (2008)
<i>Cricotopus caducus</i>	Hirvenoja, 1973		Skidmore (2008) cites Langford & Ruse (2005) first British record
<i>Cricotopus fuscus</i>	(Kieffer, 1909)		Skidmore (2008)
<i>Cricotopus obnixus</i>	(Walker, 1856)		Bland (1994)
<i>Cricotopus sylvestris</i>	(Fabricius, 1794)		Skidmore (2008)
<i>Cricotopus tricinctus</i>	(Meigen, 1818)		Skidmore (2008) cites Bland & Rotherhay (1994)
<i>Cryptochironomus supplicans</i>	(Meigen, 1830)		Waterston & Lyster (1979), Skidmore (2008)
<i>Demeijerea rufipes</i>	(Linnaeus, 1761)		Skidmore (2008)
<i>Demicryptochironomus vulneratus</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Diamesa tonsa</i>	(Haliday in Walker, 1856)		Skidmore (2008) St Kilda as <i>Diamesa culicoides</i>
<i>Endochironomus albipennis</i>	(Meigen, 1830)		Skidmore (2008)
<i>Glyptotendipes glaucus</i>	(Meigen, 1818)		Skidmore (2008)

Chironomoidea: Chironomidae - Non-biting Midges				
<i>Glyptotendipes pallens</i>	(Meigen, 1804)	<b>Requires review</b>		Waterston & Lyster (1979) as <i>G. pallens</i> f <i>glaucus</i> , Waterston (1981). Not listed by Skidmore (2008)
<i>Glyptotendipes viridis</i>	(Macquart, 1834)	<b>Requires review</b>		Unchecked by Skidmore (2008) as at the time <i>G. viridis</i> synonym of <i>G. imbecillis</i> , but <i>viridis</i> is now listed as a species
<i>Halocladus variabilis</i>	(Staeger, 1839)			Skidmore (2008) cites Buckland 1988 (archaeological dig)
<i>Halocladus varians</i>	(Staeger, 1839)			Skidmore (2008)
<i>Heterotrissocladius marcidus</i>	(Walker, 1856)			Skidmore (2008)
<i>Macropelopia adauca</i>	Kieffer in Thienemann & Kieffer, 1916			Skidmore (2008)
<i>Macropelopia nebulosa</i>	(Meigen, 1804)			Skidmore (2008)
<i>Macropelopia notata</i>	(Meigen, 1818)			Skidmore (2008)
<i>Metriocnemus eurynotus</i>	(Holmgren, 1883)			Skidmore (2008)
<i>Metriocnemus fuscipes</i>	(Meigen, 1818)			Waterston & Lyster (1979), Skidmore (2008)
<i>Metriocnemus picipes</i>	(Meigen, 1818)			Skidmore (2008)
<i>Micropsectra atrofasciata</i>	(Kieffer, 1911)			Skidmore (2008)
<i>Microtendipes chloris</i>	(Meigen, 1818)			Waterston & Lyster (1979), Skidmore (2008)
<i>Orthocladus frigidus</i>	(Zetterstedt, 1838)	St Kilda only		Skidmore (2008)
<i>Orthocladus obumbratus</i>	Kieffer, 1911	St Kilda only		Skidmore (2008), Waterston (1981) as <i>Orthocladus rhyacobius</i>
<i>Parachironomus arcuatus</i>	(Goetghebuer, 1919)			Waterston & Lyster (1979), Skidmore (2008)
<i>Parachironomus biannulatus</i>	(Staeger, 1839)			Waterston & Lyster (1979), Skidmore (2008)
<i>Parachironomus vitiosus</i>	(Goetghebuer, 1921)			Waterston & Lyster (1979), Skidmore (2008)
<i>Paramerina cingulata</i>	(Walker, 1856)			Skidmore (2008)
<i>Paratanytarsus tenuis</i>	(Meigen, 1830)			Skidmore (2008)
<i>Paratrichocladus rufiventris</i>	(Meigen, 1830)	St Kilda only		Skidmore (2008)
<i>Phaenopsectra flavipes</i>	(Meigen, 1818)			Skidmore (2008)
<i>Polypedilum albicorne</i>	(Meigen, 1838)			Skidmore (2008)
<i>Polypedilum arundineti</i>	(Goetghebuer, 1921)			Waterston & Lyster (1979), Skidmore (2008)
<i>Procladius choreus</i>	(Meigen, 1804)			Skidmore (2008)
<i>Procladius crassinervis</i>	(Zetterstedt, [1838])			Waterston & Lyster (1979), Skidmore (2008)
<i>Procladius culiciformis</i>	(Linnaeus, 1767)			Waterston & Lyster (1979), Skidmore (2008)

<b>Chironomoidea: Chironomidae - Non-biting Midges</b>			
<i>Procladius sagittalis</i>	(Kieffer, 1909)		Skidmore (2008) cites Bratton 1987
<i>Psectrocladius sordidellus</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Pseudochironomus prasinatus</i>	(Staeger, 1839)		Skidmore (2008)
<i>Pseudorthocladius filiformis</i>	(Kieffer in Kieffer & Thienemann, 1908)		Waterston (1981) as <i>Psectrocladius turfaceus</i> , Skidmore (2008),
<i>Rheotanytarsus</i>	Thienemann & Bause, 1913		Waterston & Lyster (1979), Skidmore (2008)
<i>Smittia aterrima</i>	(Meigen, 1818)	St Kilda only	Skidmore (2008)
<i>Smittia pratorum</i>	(Goetghebuer, 1927)		Skidmore (2008)
<i>Tanypus</i>	Meigen, 1803	OHBR 2017 first record	NBN: OHBR 2017 - larva
<i>Telmatogeton japonicus</i>	Tokunaga, 1933	OHBR 2010 first record	NBN: OHBR 2010
<i>Trissopelopia longimana</i>	(Staeger, 1839)		Waterston & Lyster (1979), Skidmore (2008)
<b>Tabanoidea: Rhagionidae - Snipeflies</b>			
<i>Chrysopilus cristatus</i>	(Fabricius, 1775)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2997, 1999, 2004; NBN: OHBR 2008-2017, 2019
<i>Rhagio notatus</i>	(Meigen, 1820)		Skidmore (2008)
<i>Rhagio scolopaceus</i>	(Linnaeus, 1758)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2009-2019
Records of <i>R.annulatus</i> , <i>R. scolopaceus</i> , <i>R.strigosus</i> , <i>R.tringarius</i> , <i>R. vitripennis</i> reported from Monach Islands by Majer (1988) in error			
<b>Tabanoidea: Tabanidae - Horseflies</b>			
<i>Chrysops caecutiens</i>	(Linnaeus, 1758)		Skidmore (2008) doubtful record St Kilda; NBN: OHBR 2018
<i>Chrysops relictus</i>	Meigen, 1820		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2007; NBN: BRC 1946, OHBR 2008-2019
<i>Haematopota crassicornis</i>	Wahlberg, 1848		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1999; NBN: BRC 1945, OHBR 2017
<i>Haematopota pluvialis</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1997, 1999, 2007; NBN: BRC 1945, 1960, OHBR 2008-2019
<i>Hybomitra montana</i>	(Meigen, 1820)		Forrest, Waterston & Watson (1936) as <i>Theriopectes montanus</i> , Whitely, Garland, Hancock (1994), Skidmore (2008)
<b>Stratiomyoidea: Stratiomyidae - Soldierflies</b>			
<i>Beris vallata</i>	(Forster, 1771)		Skidmore (2008); NBN: OHBR 2012-2017

Stratiomyoidea: Stratiomyidae - Soldierflies			
<i>Chloromyia formosa</i>	(Scopoli, 1763)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995, Parker 1997; NBN: OHBR 2007-2018
<i>Nemotelus nigrinus</i>	Fallén, 1817		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Nemotelus uliginosus</i>	(Linnaeus, 1767)		Skidmore (2008), Bratton (2012); NBN: OHBR 2012-2014
Asiloidea: Therevidae - Stiletto Flies			
<i>Acrosathe annulata</i>	(Fabricius, 1805)		Skidmore (2008)
<i>Thereva bipunctata</i>	Meigen, 1820		Skidmore (2008); NBN: OHBR 2012
<i>Thereva nobilitata</i>	(Fabricius, 1775)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2007; NBN: OHBR 2013
<i>Thereva plebeja</i>	(Linnaeus, 1758)	<b>Requires confirmation</b>	Skidmore (2008) cites C. W Dale - probably error for <i>T. nobilitata</i> as <i>T. plebeja</i> southern distribution. Included by Waterston (1981) as <i>T. plebeia</i> , 19th century record from North Uist - presumably C W Dale.
Empidoidea: Hybotidae - Dance Flies			
<i>Bicellaria pilosa</i>	Lundbeck, 1910		Skidmore (2008)
<i>Bicellaria spuria</i>	(Fallén, 1816)	<b>Requires review</b>	Not listed by Waterston (1981), Skidmore considers old records for Lewis and South Uist to be either <i>B. sulcata</i> or <i>B. vana</i>
<i>Chersodromia incana</i>	Haliday in Walker, 1851	OHBR 2013 first record	OHBR 2013 Mingulay
<i>Chersodromia arenaria</i>	(Haliday, 1833)	Single record St Kilda	Skidmore (2008)
<i>Chersodromia hirta</i>	(Walker, 1835)		Skidmore (1994), Skidmore (2008)
<i>Crossopalpus curvipes</i>	(Meigen, 1822)		Skidmore (1994) as <i>Drapetis curvipes</i> , Skidmore (2008)
<i>Hybos culiciformis</i>	(Fabricius, 1775)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Hybos femoratus</i>	(Müller, 1776)		Forrest, Waterston & Watson (1936), Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Platypalpus annulatus</i>	(Fallén, 1815)		Forrest, Waterston & Watson (1936) as <i>Tachydromia fulvipes</i> , Skidmore (2008)
<i>Platypalpus ciliaris</i>	(Collin, 1926)		MacGowan (1994) described as first record, Skidmore (2008)
<i>Platypalpus longicornis</i>	(Meigen, 1822)		Skidmore (1994), Skidmore 2008
<i>Platypalpus minutus</i>	(Meigen, 1804)		Skidmore (2008)
<i>Platypalpus nigritarsis</i>	(Fallén, 1816)		Skidmore (1994), Skidmore (2008)
<i>Platypalpus notatus</i>	(Meigen, 1822)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Platypalpus pallidiventris</i>	(Meigen, 1822)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay

Empidoidea: Hybotidae - Dance Flies			
<i>Platypalpus pseudofulvipes</i>	(Frey, 1909)		MacGowan (1994) as <i>P. coarctatus</i> described as first record, Skidmore (2008)
<i>Platypalpus strigifrons</i>	(Zetterstedt, 1849)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Tachydromia aemula</i>	(Loew, 1864)		Skidmore (1994), Skidmore (2008)
<i>Tachypeza fuscipennis</i>	(Fallén, 1815)		Skidmore (2008)
<i>Tachypeza nubila</i>	(Meigen, 1804)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
Empidoidea: Empididae - Dance Flies			
<i>Trichina clavipes</i>	Meigen, 1830		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Chelifera diversicauda</i>	Collin, 1927		Skidmore (1994), Skidmore (2008)
<i>Chelifera monostigma</i>	(Meigen, 1822)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Chelifera preclatoria</i>	(Fallén, 1816)		Skidmore (1994), Skidmore (2008)
<i>Clinocera bipunctata</i>	(Haliday, 1833)	OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<i>Clinocera fontinalis</i>	(Haliday, 1833)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Clinocera stagnalis</i>	(Haliday, 1833)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Empis borealis</i>	Linnaeus, 1758		Skidmore (2008); NBN: OHBR 2013
<i>Empis livida</i>	Linnaeus, 1758		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2015
<i>Empis lucida</i>	Zetterstedt, 1838		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Empis stercorea</i>	Linnaeus, 1761		Skidmore (2008); NBN: OHBR 2013, 2017
<i>Empis tessellata</i>	Fabricius, 1794		Skidmore (2008)
<i>Empis verralli</i>	Collin, 1927		Skidmore (2008) cites Roper 1995: NBN; OHBR 2013
<i>Hemerodromia baetica</i>	Collin, 1927	<b>Requires confirmation</b>	MacGowan (1994) described as first record. Not listed by Skidmore (2008)
<i>Hemerodromia raptorica</i>	Meigen, 1830		Skidmore (2008)
<i>Hilara cantabrica</i>	Strobl, 1899	NBN: <i>Hilara lundbecki</i>	Skidmore (1994), MacGowan (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Hilara choric</i>	(Fallén, 1816)		Forrest, Waterston & Watson (1936), Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Hilara clavipes</i>	(Harris, 1780)		MacGowan (1994) as <i>H. matrona</i> , Skidmore (2008)

Empidoidea: Empididae - Dance Flies			
<i>Hilara interstincta</i>	(Fallén, 1816)		MacGowan (1994) described as first record, Skidmore (2008)
<i>Hilara longivittata</i>	Zetterstedt, 1842		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Hilara maura</i>	(Fabricius, 1776)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Phyllodromia melanocephala</i>	(Fabricius, 1794)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Rhamphomyia hybotina</i>	(Zetterstedt, [1838])		Skidmore (1994), Skidmore (2008)
<i>Rhamphomyia morio</i>	Zetterstedt, 1838		Skidmore (2008)
<i>Rhamphomyia nitidula</i>	Zetterstedt, 1842		Skidmore (2008)
<i>Rhamphomyia pilifer</i>	Meigen, 1838	<b>Requires confirmation</b>	MacGowan (1994) as <i>R. dentipes</i> described as first record. Not listed by Skidmore (2008)
<i>Rhamphomyia sciarina</i>	(Fallén, 1816)		MacGowan (1994) described as first record, Skidmore (1994), Skidmore (2008)
<i>Rhamphomyia stigmosa</i>	Macquart, 1827		MacGowan (1994) described as first record, Skidmore (2008)
<i>Rhamphomyia tibiella</i>	Meigen, 1822		MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Rhamphomyia variabilis</i>	(Fallén, 1816)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013, 2018
Empidoidea: Dolichopodidae - Long-legged Flies			
<i>Achalcus flavicollis</i>	(Meigen, 1824)		Skidmore (2008)
<i>Aphrosylus celtiber</i>	Haliday, 1855		MacGowan 1987, MacGowan (1994) described as first record, Skidmore (2008); NBN: OHBR 2009-2017
<i>Aphrosylus ferox</i>	Haliday in Walker, 1851		MacGowan (1987), MacGowan (1994) as first record, Skidmore (2008); NBN: OHBR 2013
<i>Aphrosylus raptor</i>	Haliday in Walker, 1851		Skidmore (2008)
<i>Argyra argentina</i>	(Meigen, 1824)		Skidmore (2008); NBN: OHBR 2012-2013
<i>Argyra argyria</i>	(Meigen, 1824)		Skidmore (2008)
<i>Argyra diaphana</i>	(Fabricius, 1775)		MacGowan (1987), Skidmore (2008)
<i>Argyra perplexa</i>	Becker, 1918		Skidmore (2008)
<i>Argyra vestita</i>	(Wiedemann, 1817)		Skidmore (2008)
<i>Campsicnemus alpinus</i>	(Haliday, 1833)		Skidmore (2008)
<i>Campsicnemus armatus</i>	(Zetterstedt, 1849)		MacGowan (1994) described as first record, Skidmore (1994), Skidmore (2008)
<i>Campsicnemus curvipes</i>	(Fallén, 1823)		Skidmore (1994), Skidmore (2008)
<i>Campsicnemus loripes</i>	(Haliday, 1832)		Forrest, Waterston & Watson (1936), MacGowan (1987), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Campsicnemus scambus</i>	(Fallén, 1823)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: OHBR 2013
<i>Chrysotus cilipes</i>	Meigen, 1824	OHBR 2013 first record	NBN: OHBR 2013 - Mingulay



Empidoidea: Dolichopodidae - Long-legged Flies			
<i>Chrysotus gramineus</i>	(Fallén, 1823)		Skidmore (2008)
<i>Dolichopus acuticornis</i>	Wiedemann, 1817		Forrest, Waterston & Watson (1936), MacGowan (1987), Skidmore (2008) cites Roper 1995
<i>Dolichopus atratus</i>	Meigen, 1824		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2010-2013
<i>Dolichopus atripes</i>	Meigen, 1824		Forrest, Waterston & Watson (1936), MacGowan (1987), Skidmore (1994), Skidmore (2008)
<i>Dolichopus brevipennis</i>	Meigen, 1824		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Dolichopus caligatus</i>	Wahlberg, 1850		MacGowan (1987), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Dolichopus diadema</i>	Haliday, 1832	St Kilda records only	Waterston (1981) as <i>Macrodolichopus diadema</i> , Skidmore (2008)
<i>Dolichopus festivus</i>	Haliday, 1832	OHBR 2013 first record	NBN: OHBR 2013
<i>Dolichopus griseipennis</i>	Stannius, 1831		Forrest, Waterston & Watson (1936), Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013
<i>Dolichopus lepidus</i>	Staeger, 1842		MacGowan (1994) as first record Lewis, Skidmore (2008)
<i>Dolichopus longicornis</i>	Stannius, 1831		Skidmore (2008)
<i>Dolichopus longitarsis</i>	Stannius, 1832		MacGowan (1994) Flannan Islands, Skidmore (2008)
<i>Dolichopus nubilus</i>	Meigen, 1824		MacGowan (1987), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Bacchus 1966 Monach Islands; NBN: OHBR 2012-2013
<i>Dolichopus pennatus</i>	Meigen, 1824		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013
<i>Dolichopus plumipes</i>	(Scopoli, 1763)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2010-2013
<i>Dolichopus popularis</i>	Wiedemann, 1817		MacGowan (1987), MacGowan (1994) described as first record, Skidmore (2008)
<i>Dolichopus rupestris</i>	Haliday, 1833		MacGowan (1987), Skidmore (2008)
<i>Dolichopus signifer</i>	Haliday, 1838	<b>Requires review</b>	Skidmore (2008) - doubtful. See MacGowan (1986, 1994). Dale (1884) records <i>D. punctum</i> which Grimshaw (1914) considered doubtful
<i>Dolichopus simplex</i>	Meigen, 1824		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2012
<i>Dolichopus trivialis</i>	Haliday, 1832		Forrest, Waterston & Watson (1936), Skidmore (1994), MacGowan (1987), Skidmore (2008)
<i>Dolichopus unguatus</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012
<i>Dolichopus urbanus</i>	Meigen, 1824		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Dolichopus vitripennis</i>	Meigen, 1824		Skidmore (2008) cites Roper 1995
<i>Gymnopternus aerosus</i>	(Fallén, 1823)	NBN: <i>Hercostomus aerosus</i>	Forrest, Waterston & Watson (1936) as <i>Hercostomus aerosus</i> , Skidmore (2008)

Empidoidea: Dolichopodidae - Long-legged Flies				
<i>Gymnopternus cupreus</i>	(Fallén, 1823)	NBN: <i>Hercostomus cupreus</i>		Skidmore (2008) - Vardy & Bromfield 1962 Stornoway.
<i>Hercostomus germanus</i>	(Wiedemann, 1817)	OHBR 2012 first record		NBN: OHBR 2012
<i>Hydrophorus albiceps</i>	Frey, 1915			MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Hydrophorus balticus</i>	(Meigen, 1824)			Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (1994)
<i>Hydrophorus nebulosus</i>	(Fallén, 1823)			Skidmore (2008)
<i>Hydrophorus praecox</i>	(Lehmann, 1822)			Skidmore (2008)
<i>Liancalus virens</i>	(Scopoli, 1763)			Forrest, Waterston & Watson (1936), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Medetera jugalis</i>	Collin, 1941	<b>Requires review</b>		“sparse records from Highland and the Western Isles” (Drake 2018). McGowan notes very rare in Scotland. Skidmore (2008) cites a record from Skye from 1990
<i>Medetera petrophiloides</i>	Parent, 1925			Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Medetera truncorum</i>	Meigen, 1824			MacGowan (1994) described as first record, Skidmore (2008)
<i>Micromorphus albipes</i>	(Zetterstedt, 1843)			Skidmore (2008)
<i>Microphor crassipes</i>	Macquart, 1827			Skidmore (2008) - Collected by C.W. Dale in St Kilda, specimen Oxford Museum (Davies 1981)
<i>Rhaphium brevicorne</i>	Curtis, 1835	OHBR 2013 first record		NBN: OHBR 2013 Mingulay
<i>Rhaphium consobrinum</i>	Zetterstedt, 1843			MacGowan (1987), MacGowan (1994) described as first record, Skidmore (2008)
<i>Rhaphium crassipes</i>	(Meigen, 1824)			MacGowan (1994) described as first record Lewis, Skidmore (2008)
<i>Rhaphium monotrichum</i>	Loew, 1850			Skidmore (2008)
<i>Sciapus contristans</i>	(Wiedemann, 1817)	<b>Not confirmed</b>		Skidmore (2008) -Grimshaw (1914) probably <i>S. zonatulus</i> . MacGowan (1987) records later corrected to <i>S. zonatulus</i>
<i>Sciapus zonatulus</i>	(Zetterstedt, 1843)			MacGowan (1987) originally recorded as <i>S. contristans</i> , MacGowan (1994), Skidmore (2008)
<i>Sympycnus cirripes</i>	(Haliday in Walker, 1851)			MacGowan (1987), Skidmore (1994), Skidmore 2008
<i>Sympycnus pulicarius</i>	(Fallén, 1823)			Forrest, Waterston & Watson (1936) as <i>S. annulipes</i> , later as <i>S. desoutteri</i> by Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012-2013
<i>Syntormon denticulatus</i>	(Zetterstedt, 1843)			As <i>S. denticulatum</i> MacGowan (1994) described as first record, Skidmore (2008)
<i>Syntormon monilis</i>	(Fabricius, 1794)			As <i>S. monile</i> Skidmore (2008)

<b>Empidoidea: Dolichopodidae - Long-legged Flies</b>			
<i>Syntormon pallipes</i>	(Fabricius, 1794)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012
<i>Syntormon pumilus</i>	(Meigen, 1824)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Syntormon tarsatum</i>	(Fallén, 1823)		Forrest, Waterston & Watson (1936), MacGowan (1987), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Tachytrechus notatus</i>	(Stannius, 1831)		Skidmore (2008)
<i>Thrypticus bellus</i>	Loew, 1869		MacGowan (1994) described as first record, Skidmore (2008)
<i>Thrypticus divisus</i>	(Strobl, 1880)		MacGowan (1994) described as first record, Skidmore (2008) quotes Falk and Crossley (2005) who cite distribution in Scotland to be Easternness and Western Isles
<b>Platypezoidea: Phoridae - Scuttle Flies</b>			
<i>Phora stictica</i>	Meigen, 1830		Skidmore (2008) cites Barron 1987
<b>Lonchopteroidea: Lonchopteridae - Pointed-wing Flies</b>			
<i>Lonchoptera bifurcata</i>	(Fallén, 1823)		Skidmore (1994), Skidmore (2008)
<i>Lonchoptera lutea</i>	Panzer, 1809		MacGowan (1994) Flannan Islands, Skidmore (1994), Skidmore (2008) cites Roper 1995, Buckland 1988; NBN: OHBR 2013 Mingulay
<b>Syrphoidea: Syrphidae - Hoverflies</b>			
<i>Anasimyia lineata</i>	(Fabricius, 1787)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997; NBN: OHBR 2018
<i>Anasimyia lunulata</i>	(Meigen, 1822)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1999
<i>Cheilosia albitarsis</i>	(Meigen, 1822)	OHBR 2013 first record	NBN: OHBR 2013
<i>Cheilosia bergenstammi</i>	Becker, 1894		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997, 1999; NBN: OHBR 1998-2017
<i>Cheilosia fraterna</i>	(Meigen, 1830)	OHBR 2013 first record	NBN: OHBR 2013
<i>Cheilosia illustrata</i>	(Harris, [1780])		Forrest, Waterston & Watson (1936), MacGowan (1987), Skidmore (2008) cites Bratton 1987, Parker 1997, 1999; NBN: OHBR 1998-2019
<i>Cheilosia latifrons</i>	(Zetterstedt, 1843)		Skidmore (2008) cites Parker 2007
<i>Cheilosia pagana</i>	(Meigen, 1822)		Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Chrysotoxum arcuatum</i>	(Linnaeus, 1758)		Skidmore (2008)
<i>Chrysotoxum bicinctum</i>	(Linnaeus, 1758)	OHBR 2014 first record	NBN: OHBR 2014
<i>Dasysyrphus albostriatus</i>	(Fallén, 1817)	OHBR 2012 first record	NBN: OHBR 2012
<i>Dasysyrphus pinastri</i>	(De Geer, 1776)	<b>Requires review</b>	Skidmore (2008) cites a record for Lewis and equates <i>D. pinastri</i> with <i>D. lunulatus</i> (Meigen). It is unclear whether this is MacGowan's record

Syrphoidea: Syrphidae - Hoverflies			
<i>Dasysyrphus venustus</i>	(Meigen, 1822)	Requires review	MacGowan (1994) as <i>D. lunulatus</i> Meig described as first record, which is now a synonym of <i>D. venustus</i> which is not listed by Skidmore (2008)
<i>Episyrphus balteatus</i>	(De Geer, 1776)		Whitely, Garland, Hancock (1994), Skidmore (1994), MacGowan (1987), Skidmore (2008) cites Parker 1999; NBN: OHBR 2007-2019
<i>Eristalinus aeneus</i>	(Scopoli, 1763)		Whitely, Garland, Hancock (1994), Bratton (2012), Skidmore (2008) cites Roper 1995 Parker 2007; NBN: OHBR 2010-2013
<i>Eristalinus sepulchralis</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936) as <i>Eristalis sepulchalis</i> , Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997, 1999, Roper 1995; NBN: OHBR 2007-2013
<i>Eristalis abusivus</i>	Collin, 1931		Skidmore (1994), Skidmore (2008) cites Parker 1999, 2005; NBN: OHBR 2010-2013
<i>Eristalis arbustorum</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1997-2007; NBN: OHBR 2009-2019
<i>Eristalis horticola</i>	(De Geer, 1776)		Forrest, Waterston & Watson (1936), Skidmore (2008) cites Parker 2006; NBN: OHBR 2009-2018
<i>Eristalis intricarius</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), MacGowan (1994), Skidmore (2008) cites Smith 1955, Roper 1995. Parker 1997-2007; NBN: OHBR 2002-2019
<i>Eristalis nemorum</i>	(Linnaeus, 1758)		Skidmore (2008) cites Parker 1999; NBN: OHBR 2009-2017
<i>Eristalis pertinax</i>	(Scopoli, 1763)		Skidmore (2008) cites Roper 1995, Parker 1997, 2005; NBN: OHBR 2001-2019
<i>Eristalis tenax</i>	(Linnaeus, 1758)		Skidmore (2008) cites Parker 2002; NBN: OHBR 2010-2017
<i>Eupeodes corollae</i>	(Fabricius, 1794)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1999-2007; NBN: OHBR 2009-2017, 2019
<i>Eupeodes latifasciatus</i>	(Macquart, 1829)		MacGowan (1994), Skidmore (2008) cites Parker 2004-2007; NBN: OHBR 2008-2017
<i>Eupeodes lundbecki</i>	(Loew, 1843)	Rare vagrant (Ball & Morris 2013)	Skidmore (2008)
<i>Eupeodes luniger</i>	(Meigen, 1822)		Skidmore (2008) cites Parker 1997, 1999; NBN: OHBR 2009
<i>Helophilus hybridus</i>	Loew, 1846		Skidmore (1994), Skidmore (2008), NBN; OHBR 2013-2018
<i>Helophilus pendulus</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997 - 2007; NBN: OHBR 2008-2019
<i>Helophilus trivittatus</i>	(Fabricius, 1805)	OHBR 2008 first record	NBN: OHBR 2008, 2013
<i>Lejogaster metallina</i>	(Fabricius, 1781)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1999-2006; NBN: OHBR 2008-2017
<i>Leucozona glaucia</i>	(Linnaeus, 1758)		Skidmore (2008); NBN: OHBR 2011-2017, 2019
<i>Leucozona lucorum</i>	(Linnaeus, 1758)	OHBR 2011 first record	NBN: OHBR 2011-2018

Syrphoidea: Syrphidae - Hoverflies			
<i>Melangyna lasiophthalma</i>	(Zetterstedt, 1843)	OHBR 2013 first record	NBN: OHBR 2013
<i>Melanogaster aerosa</i>	(Loew, 1843)		MacGowan (1994) described as first record, Skidmore (1994), Skidmore (2008)
<i>Melanogaster hirtella</i>	(Loew, 1843)		Forrest, Waterston & Watson (1936) as <i>Chrysogaster hirtella</i> , Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995, Parker 1997-2002; NBN: OHBR 2013-2017
<i>Melanostoma dubium</i>	(Zetterstedt, 1837)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008). The genus <i>Melanostoma</i> is the subject of current investigation and the concept of species recognised here may change. The status of <i>dubium</i> still needs to be resolved Chandler (2019)
<i>Melanostoma mellinum</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1997- 200; NBN: OHBR 2008-2017
<i>Melanostoma scalare</i>	(Fabricius, 1794)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1997- 2007; NBN: OHBR 2009-2019
<i>Meliscaeva auricollis</i>	(Meigen, 1822)		Skidmore (2008); NBN: OHBR 2002, 2012, 2019
<i>Meliscaeva cinctella</i>	(Zetterstedt, 1843)	OHBR 2012 first record	NBN: OHBR 2012, 2013, 2019
<i>Neoascia podagrica</i>	(Fabricius, 1775)		Forrest, Waterston & Watson (1936), Skidmore (2008) cites Parker 2006, 2007; NBN: OHBR 2013, 2014, 2019
<i>Neoascia tenur</i>	(Harris, [1780])		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1997, 1999, 2004, 2007; NBN: OHBR 2009, 2013
<i>Parasyrphus punctulatus</i>	(Verrall, 1873)		MacGowan (1994) described as first record, Skidmore (2008)
<i>Pipizella viduata</i>	(Linnaeus, 1758)		Skidmore (2008) as <i>P. virens</i> but presumed to be to be <i>P. viduata</i> ; NBN: OHBR 2013
<i>Platycheirus albimanus</i>	(Fabricius, 1781)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997-2007; NBN: OHBR 2009-2017
<i>Platycheirus angustatus</i>	(Zetterstedt, 1843)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997 - 2007; NBN: OHBR 2012
<i>Platycheirus clypeatus</i>	(Meigen, 1822)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), MacGowan (1987), Skidmore (2008) cites Roper 1995, Parker 1997-2007; NBN: OHBR 2011-2013
<i>Platycheirus granditarsus</i>	(Forster, 1771)		MacGowan (1987), MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997, 1999; NBN: OHBR 2012
<i>Platycheirus immarginatus</i>	(Zetterstedt, 1849)		Whitely, Garland, Hancock (1994), MacGowan (1987), Skidmore (2008)
<i>Platycheirus manicatus</i>	(Meigen, 1822)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Smith 1955, Roper 1995, Parker 1997-2007; NBN: OHBR 2008-2017, 2019
<i>Platycheirus melanopsis</i>	Loew, 1856	OHBR 2013 first record	NBN: OHBR 2013

Syrphoidea: Syrphidae - Hoverflies				
<i>Platycheirus nielseni</i>	Vockeroth, 1990	OHBR 2013 first record	NBN: OHBR 2013	
<i>Platycheirus occultus</i>	Goeldlin de Tiefenau Maibach & Speight, 1990		Skidmore (2008); NBN: OHBR 2013	
<i>Platycheirus peltatus</i>	(Meigen, 1822)		Skidmore (1994), Skidmore (2008)	
<i>Platycheirus podagratus</i>	(Zetterstedt, [1838])		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1999, 2007; NBN: OHBR 2013	
<i>Platycheirus ramsarensis</i>	Goeldlin de Tiefenau Maibach & Speight, 1990		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1999; NBN: OHBR 2009	
<i>Platycheirus scambus</i>	(Staeger, 1843)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), MacGowan (1987), Skidmore (2008) cites Parker 1999; NBN: OHBR 2009	
<i>Platycheirus scutatus</i>	(Meigen, 1822)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2004; NBN: OHBR 2013	
<i>Platycheirus splendidus</i>	Rotheray, 1998		Skidmore (2008) cites Parker 1999	
<i>Platycheirus sticticus</i>	(Meigen, 1822)		MacGowan (1994) described as first record, Skidmore (2008) cites Bratton 1987	
<i>Rhingia campestris</i>	Meigen, 1822		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997, 1999; NBN: OHBR 1998-2019	
<i>Scaeva pyrastris</i>	(Linnaeus, 1758)		MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2004, 2007; NBN: OHBR 2004-2019	
<i>Sericomyia lappona</i>	(Linnaeus, 1758)		Whitely, Garland, Hancock (1994) Skidmore (2008); NBN: OHBR 2014-2018	
<i>Sericomyia silentis</i>	(Harris, [1776])		Forrest, Waterston & Watson (1936) as <i>S. borealis</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1999, 2004; NBN: OHBR 1990-2019	
<i>Sphaerophoria interrupta</i>	(Fabricius, 1805)		Forrest, Waterston & Watson (1936), as <i>S. menthastris</i> var. <i>picta</i> , Skidmore (2008) cites Parker 2004, 2007; NBN: OHBR 2012-2013	
<i>Sphaerophoria scripta</i>	(Linnaeus, 1758)		Wakeham-Dawson (2007), Skidmore (2008)	
<i>Sphegina clunipes</i>	(Fallén, 1816)	OHBR 2014 first record	NBN: OHBR 2014	
<i>Syritta pipiens</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), MacGowan (1987), Skidmore (2008) cites Parker 1997, 1999, 2007; NBN: OHBR 2009-2019	
<i>Syrphus ribesii</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), MacGowan (1987), Skidmore (2008) cites Parker 2004; NBN: OHBR 2008-2018	
<i>Syrphus torvus</i>	Osten Sacken, 1875		MacGowan (1994) described as first record, Skidmore (2008); NBN: OHBR 2012, 2013	
<i>Syrphus vitripennis</i>	Meigen, 1822		Skidmore (1994), MacGowan (1987), Skidmore (2008); NBN: OHBR 2002-2018	
<i>Trichopsomyia flavitarsis</i>	(Meigen, 1822)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997; NBN: OHBR 2013	
<i>Tropidia scita</i>	(Harris, [1780])		MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013-2019	



<b>Syrphoidea: Syrphidae - Hoverflies</b>			
<i>Volucella bombylans</i>	(Linnaeus, 1758)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Smith 1955, Parker 1997, 1999, 2004; NBN: OHBR 2002-2018
<i>Volucella pellucens</i>	(Linnaeus, 1758)	OHBR 2010 first record	NBN: OHBR 2010-2017
<i>Xanthandrus comtus</i>	(Harris, 1780)		Skidmore (2008)? Not recorded in Scotland NBN. Migratory (Ball & Morris 2013)
<i>Xylota segnis</i>	(Linnaeus, 1758)	OHBR 2012 first record	NBN: OHBR 2012-2016, 2018
<b>Nerioidea: Micropezidae - Stilt-flies</b>			
<i>Calobata petronella</i>	(Linnaeus, 1761)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013, 2015
<b>Diopsoidea: Psilidae - Rust Flies</b>			
<i>Chamaepsila atra</i>	(Meigen, 1826)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<b>Diopsoidea: Psilidae - Rust Flies</b>			
<i>Chamaepsila nigra</i>	(Fallén, 1820)		Skidmore (2008) cites Roper 1995
<i>Chamaepsila nigricornis</i>	(Meigen, 1826)		Skidmore (2008); NBN: Dipterists Forum 1974
<i>Chamaepsila pallida</i>	(Fallén, 1820)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Chamaepsila rosae</i>	(Fabricius, 1794)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Parker 1999
<i>Loxocera aristata</i>	(Panzer, 1801)		Whitely, Garland, Hancock (1994) as <i>Loxocera ichneumone</i> , Skidmore (2008); NBN: Dipterists Forum 2006
<b>Tephritoidea: Pallopteridae - Trembling-wing Flies</b>			
<i>Palloptera quinquemaculata</i>	(Macquart, 1835)	OHBR 2013 first record	NBN: OHBR 2013
<i>Palloptera ustulata</i>	Fallén, 1820		Forrest, Waterston & Watson (1936), Skidmore (2008)
<b>Tephritoidea: Piophilidae - Skippers</b>			
<i>Allopiophila vulgaris</i>	(Fallén, 1820)		Waterston (1981) as <i>Piophila vulgaris</i> , Skidmore (2008)
<i>Prochyliza nigrimana</i>	(Meigen, 1826)		Skidmore (2008)
<i>Prochyliza varipes</i>	(Meigen, 1830)	NBN: <i>Liopiophila varipes</i>	Skidmore (1994), Skidmore (2008) cites Roper 1995.
<i>Stearibia nigriceps</i>	(Meigen, 1826)		Skidmore (2008) cites Roper 1995
<b>Tephritoidea: Ulidiidae - Picture-winged Flies</b>			
<i>Herina frondescentiae</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 1997; NBN: OHBR 2009-20015
<b>Tephritoidea: Platystomatidae - Signal Flies</b>			
<i>Rivellia syngenesiae</i>	(Fabricius, 1781)		Skidmore (2008) cites Parker 2007



Tephritoidea: Tephritidae - Gallflies			
<i>Campiglossa absinthii</i>	(Fabricius, 1805)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2007
<i>Campiglossa loewiana</i>	(Hendel, 1927)		MacGowan (1994) described as first record, Skidmore (2008)
<i>Chaetostomella cylindrica</i>	(Robineau-Desvoidy, 1830)		Skidmore (2008) cites Parker 2007
<i>Ensina sonchi</i>	(Linnaeus, 1767)	<b>Requires confirmation</b>	Nottingham Museum - Lewis n.d. - only UK record on NBN. Not listed by Skidmore (2008)
<i>Euleia heraclei</i>	(Linnaeus, 1758)		Bland (1994), Skidmore (2008) cites Parker 2004; NBN: OHBR 2017
<i>Sphenella marginata</i>	(Fallén, 1814)		Skidmore (2008) cites Parker 2007
<i>Trupanea stellata</i>	(Fuessly, 1775)		Skidmore (2008)
<i>Trypeta artemisiae</i>	(Fabricius, 1794)		Bland (1994), Skidmore (2008) cites Parker 2007
<i>Xyphosia miliaria</i>	(Schrank, 1781)	OHBR 2017 first record	OHBR 2017
Lauxanioidea: Lauxaniidae			
<i>Calliopum aeneum</i>	(Fallén, 1820)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Calliopum simillimum</i>	(Collin, 1933)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Calliopum tuberculatum</i>	(Becker, 1895)	NBN: <i>Sapromyza halidayi</i>	Waterston (198) as <i>Sapromyza sordida</i> , Skidmore (2008) as <i>Sapromyza halidayi</i> cites Roper 1995.
<i>Meiosimyza rorida</i>	(Fallén, 1820)		Forrest, Waterston & Watson (1936) as <i>Sapromyza rorida</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Minettia tubifer</i>	(Meigen, 1826)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Pseudolyciella stylata</i>	(Papp, 1978)	<b>Requires review</b>	Skidmore (2008) assumes all published records of <i>Lyciella palliventris</i> to be <i>P. stylata</i> . Is this an error as <i>P. palliventris</i> is recorded in Scotland, but <i>P. stylata</i> is not?
<i>Sapromyza albiceps</i>	Fallén, 1820		Whitely, Garland, Hancock (1994), Skidmore (2008)
Lauxanioidea: Chamaemyiidae			
<i>Chamaemyia flavipalpis</i>	(Haliday, 1838)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: iRecord -Whitely 1989 - presumably the same record from Eriskay?
<i>Chamaemyia geniculata</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Chamaemyia juncorum</i>	(Fallén, 1823)		Skidmore (2008)
<i>Chamaemyia polystigma</i>	(Meigen, 1830)		Skidmore (2008)
Sciomyzoidea: Coelopidae - Kelp Flies			
<i>Coelopa frigida</i>	(Fabricius, 1805)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012-2017
<i>Coelopa pilipes</i>	Haliday, 1838		Skidmore (2008); NBN: OHBR 2013-2017

Sciomyzoidea: Dryomyzidae			
<i>Neuroctena anilis</i>	(Fallén, 1820)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2018
Sciomyzoidea: Helcomyzidae			
<i>Helcomyza ustulata</i>	Curtis, 1825		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2012
Sciomyzoidea: Heterocheilidae			
<i>Heterocheila buccata</i>	(Fallén, 1820)		Forrest, Waterston & Watson (1936) as <i>Oedoparea buccata</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
Sciomyzoidea: Sciomyzidae - Snail-killing Flies			
<i>Dictya umbrarum</i>	(Linnaeus, 1758)		Skidmore (2008)
<i>Ditaeniella grisescens</i>	(Meigen, 1830)		Skidmore (2008)
<i>Hydromya dorsalis</i>	(Fabricius, 1775)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: OHBR 2012-2013
<i>Ilione albisetata</i>	(Scopoli, 1763)		Forrest, Waterston & Watson (1936) as <i>Elgiva albisetata</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Ilione lineata</i>	(Fallén, 1820)		Skidmore (1994), Skidmore (2008)
<i>Limnia paludicola</i>	Elberg, 1965		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Limnia unguicornis</i>	(Scopoli, 1763)		MacGowan (1994) described as first record, Skidmore (2008)
<i>Pherbellia cinerella</i>	(Fallén, 1820)		Forrest, Waterston & Watson (1936) as <i>Ditienia cinerella</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Pherbina coryleti</i>	(Scopoli, 1763)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Pteromicra angustipennis</i>	(Staeger, 1845)		Skidmore (2008)
<i>Renocera strobli</i>	Hendel, 1900		MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Tetanocera arrogans</i>	Meigen, 1830		Skidmore (2008)
<i>Tetanocera elata</i>	(Fabricius, 1781)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2012
<i>Tetanocera ferruginea</i>	Fallén, 1820		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Tetanocera fuscinervis</i>	(Zetterstedt, 1838)		MacGowan (1994) described as first record, Whitely, Garland, Hancock (1994), Skidmore (2008)

\*BRC records on NBN of Sepsidae from Britain and Ireland to 1985 are based on identification of museum specimens by Pont (1986), but the name of the collector is not stated. However, from the dates and locations some of these records can be identified from the published literature and are included in Waterston (1981) and Skidmore (2008)

<b>Sciomyzoidea: Sciomyzidae - Snail-killing Flies</b>			
<i>Tetanocera hyalipennis</i>	von Roser, 1840		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Tetanocera robusta</i>	Loew, 1847		Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Trypetoptera punctulata</i>	(Scopoli, 1763)	OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<b>Sciomyzoidea: Sepsidae - Ensign Flies</b>			
<i>Nemopoda nitidula</i>	Fallén, 1820		Forrest, Waterston & Watson (1936) as <i>N. cylindrica</i> , Skidmore (1994), Skidmore (2008), NBN: BRC 1905-1976*
<i>Orygma luctuosum</i>	Meigen, 1830		Skidmore (1994), Skidmore (2008), cites Roper 1995; NBN: OHBR 2014-2017
<i>Sepsis cynipsea</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: BRC 1906-1966*, OHBR 2013
<i>Sepsis flavimana</i>	Meigen, 1826		Skidmore (1994), Skidmore (2008); NBN: BRC 1973, 1976*
<i>Sepsis fulgens</i>	Meigen, 1826		Skidmore (1994), Skidmore (2008); NBN: BRC 1935-1936*, OHBR 2013
<i>Sepsis orthocnemis</i>	Frey, 1908		Skidmore (1994), Skidmore (2008); NBN: BRC 1906-1966*, OHBR 2013,
<i>Sepsis punctum</i>	(Fabricius, 1794)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2019
<i>Themira annulipes</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2017
<i>Themira leachi</i>	(Meigen, 1826)		Skidmore (2008); NBN: BRC 1962*
<i>Themira lucida</i>	(Staeger in Schjødtte, 1844)		Skidmore (1994), Skidmore (2008); NBN: BRC 1906*
<i>Themira minor</i>	(Haliday, 1833)		Skidmore (1994), Skidmore (2008)
<i>Themira pusilla</i>	(Zetterstedt, 1847)		Skidmore (1994), Skidmore (2008)
<i>Themira putris</i>	(Linnaeus, 1758)		Skidmore (2008); NBN: BRC 1905*
<i>Themira superba</i>	(Haliday, 1833)		Skidmore (1994), Skidmore (2008)
<b>Opomyzoidea: Agromyzidae - Leaf-miner Flies</b>			
<i>Agromyza mobilis</i>	Meigen, 1830		Skidmore (2008) cites Bratton 1987
<i>Agromyza nana</i>	Meigen, 1830		Bland (1994); Skidmore (2008); NBN: OHBR 2018
<i>Agromyza nigripes</i>	Meigen, 1831	St Kilda record(s) only	Skidmore (2008)
<i>Aulagromyza hendeliana</i>	(Hering, 1926)	OHBR 2017 first record	NBN: OHBR 2017
<i>Cerodontha denticornis</i>	(Panzer, [1806])		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Broomfield & Vardy 1962
<i>Cerodontha iraeos</i>	(Robineau-Desvoidy, 1851)		Bland (1994), Skidmore (2008)
<i>Chromatomyia aprilina</i>	(Goureau, 1851)	OHBR 2018 first record	NBN: OHBR 2018, 2019
<i>Chromatomyia asteris</i>	(Hendel, 1934)		As <i>Phytomyza asteris</i> , Bland (1994), Skidmore (2008)

Opomyzoidea: Agromyzidae - Leaf-miner Flies				
<i>Chromatomyia fuscula</i>	(Zetterstedt, 1838)	St Kilda record(s) only		Skidmore (2008)
<i>Chromatomyia horticola</i>	(Goureau, 1851)			As <i>Phytomyza horticola</i> , Bland (1994), Skidmore (2008)
<i>Chromatomyia loniceræ</i>	(Robineau-Desvoidy, 1851)		OHBR 2015 first record	NBN: OHBR, 2015-2018
<i>Chromatomyia primulae</i>	(Robineau-Desvoidy, 1851)		OHBR 2018 first record	NBN: OHBR 2018
<i>Chromatomyia syngenesiae</i>	Hardy, 1849			Skidmore (2008); NBN: OHBR 2018
<i>Liriomyza flaveola</i>	(Fallén, 1823)			Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Liriomyza lutea</i>	Meigen, 1830			Skidmore (2008)
<i>Liriomyza richteri</i>	Hering, 1927			Skidmore (1994), Skidmore (2008)
<i>Napomyza lateralis</i>	(Fallén, 1823)			Skidmore (2008)
<i>Phytomyza affinis</i>	(Fallén, 1823)			Skidmore (2008) - Flannan Islands - unpublished record
<i>Phytomyza alpina</i>	Groschke, 1957			Bland (1994)
<i>Phytomyza angelicae</i>	Kaltenbach, 1872			Bland (1994), Skidmore (2008); NBN: OHBR 2017, 2018
<i>Phytomyza calthophila</i>	Hendel, 1931			Bland (1994); NBN: OHBR 2016, 2019
<i>Phytomyza ilicis</i>	Curtis, 1846		OHBR 2015 first record	NBN: OHBR 2015
<i>Phytomyza lappae</i>	Goureau, 1851			Skidmore (2008)
<i>Phytomyza plantaginis</i>	Robineau-Desvoidy, 1851			Bland (1994)
<i>Phytomyza ranunculi</i>	(Schrank, 1803)			Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008); NBN: OHBR 2018, 2019
<i>Phytomyza spondylii</i>	Robineau-Desvoidy, 1851			Bland (1994); NBN: OHBR 2015 (as <i>Phytomyza spondylii</i> or <i>P. pastinacae</i> )
<i>Phytomyza varipes</i>	Macquart, 1835			Skidmore (2008)
<i>Phytomyza virgaureae</i>	Hering, 1926			Bland (1994)
<i>Phytomyza wahlgreni</i>	Rydén, 1944			Skidmore (2008) cites Bratton 1987, Skidmore 1989
Opomyzoidea: Opomyzidae				
<i>Geomyza balachowskyi</i>	Mesnil, 1934			Forrest, Waterston & Watson (1936) as <i>G. combinata</i> , Skidmore (1994), Skidmore (2008)
<i>Geomyza tripunctata</i>	Fallén, 1823			Skidmore (2008) cites Roper 1995
<i>Geomyza venusta</i>	(Meigen, 1830)			Skidmore (2008)
<i>Opomyza germinationis</i>	(Linnaeus, 1758)			Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Bratton 1987; NBN: OHBR 2013 Mingulay

Opomyzoidea: Opomyzidae			
<i>Opomyza petrei</i>	Mesnil, 1934		Skidmore (1994), Skidmore (2008)
Opomyzoidea: Anthomyzidae			
<i>Anthomyza gracilis</i>	Fallén, 1823		Skidmore (2008)
<i>Anthomyza pallida</i>	(Zetterstedt, 1838)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Anthomyza paraneglecta</i>	Elberg, 1968		Skidmore (2008)
<i>Arganthomyza socculata</i>	(Zetterstedt, 1847)		Skidmore (2008) as <i>Anthomyza socculata</i>
Carnoidea: Milichiidae			
<i>Phyllomyza securicornis</i>	Fallén, 1823		Skidmore (2008)
Carnoidea: Carnidae - Bird Flies			
<i>Meoneura flavifacies</i>	Collin, 1930		Skidmore (1994), Skidmore (2008) cites Bratton 1987
Carnoidea: Carnidae - Bird Flies			
<i>Meoneura lamellata</i>	Collin, 1930		Skidmore (2008) cites Roper 1995
<i>Meoneura obscurella</i>	(Fallén, 1823)	St Kilda record(s) only	Skidmore (2008)
Carnoidea: Chloropidae - Phantom-midges			
<i>Camarota curvipennis</i>	(Latreille, 1805)	<b>Requires confirmation</b>	NBN: Nottingham City Museums - no date. Doubtful as species distribution according to NBN SE England. See Falk, Ismay, & Chandler (2016)
<i>Cetema cereis</i>	Fallén, 1820	St Kilda record(s) only	Skidmore (2008) St Kilda
<i>Cetema elongatum</i>	(Meigen, 1830)		Forrest, Waterston & Watson (1936), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Cetema neglectum</i>	Tonnoir, 1921		MacGowan (1994) described as first record, Skidmore (2008),
<i>Cetema simile</i>	Ismay, 1985	<b>Requires review</b>	Skidmore (2008) notes that early records of <i>Cetema</i> need to checked
<i>Chlorops calceatus</i>	Meigen, 1830		Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Chlorops hypostigma</i>	Meigen, 1830	St Kilda record(s) only	Skidmore (2008)
<i>Chlorops laetus</i>	Meigen, 1830		Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Chlorops meigenii</i>	Loew, 1866		Skidmore (2008)
<i>Chlorops pumilionis</i>	(Bjerkander, 1778)		Skidmore (2008)
<i>Chlorops scalaris</i>	Meigen, 1830		Skidmore (2008)
<i>Elachiptera cornuta</i>	(Fallén, 1820)		Skidmore (1994), Skidmore (2008) cites Bratton 1987
<i>Epiochlorops puncticollis</i>	(Zetterstedt, 1848)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Melanum laterale</i>	(Haliday, 1833)		Skidmore (2008)

Carnoidea: Chloropidae - Phantom-midges				
<i>Meromyza meigeni</i>	Nartshuk, 2006		<b>Requires review</b>	Skidmore (2008)? Possible error for <i>M. bohémica</i> . Recorded Barra, not listed by Waterston (1981)
<i>Meromyza saltatrix</i>	(Linnaeus, 1761)			Skidmore (2008)
<i>Meromyza pratorum</i>	Meigen, 1830			Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013 as <i>M. pratorum</i> agg. Mingulay
<i>Oscinella frit</i>	(Linnaeus, 1758)			Forrest, Waterston & Watson (1936) as <i>Oscinosoma frit</i> , Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Oscinella pusilla</i>	(Meigen, 1830)		St Kilda record(s) only	Skidmore (2008)
<i>Oscinella vastator</i>	(Curtis, 1845)			Skidmore (2008)
<i>Oscinella vindicata</i>	(Meigen, 1830)			Skidmore (1994), Skidmore (2008) cites Roper 1995
Sphaeroceroidea: Heleomyzidae				
<i>Heleomyza borealis</i>	Boheman, 1865			Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013 Mingulay
<i>Heteromyza commixta</i>	Collin, 1901			Skidmore (1994), Skidmore (2008)
<i>Neoleria inscripta</i>	(Meigen, 1830)			Skidmore (2008) cites Roper 1995
<i>Neoleria maritima</i>	(Villeneuve, 1921)			Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Neoleria prominens</i>	(Becker, 1897)			Edwards & Collin (1932), Lack (1932), Skidmore (2008), Horsfield (2013)- St Kilda, Flannan Islands
<i>Oecothea praecox</i>	Loew, 1862		OHBR 2013 first record	NBN: OHBR 2013 Mingulay
<i>Suillia bicolor</i>	(Zetterstedt, [1838])			Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997, 1999
<i>Suillia imberbis</i>	Czerny, 1924			Skidmore (2008)
<i>Suillia notata</i>	(Meigen, 1830)			Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995, Parker 1997, 1999
<i>Tephrochlamys rufiventris</i>	(Meigen, 1830)			Skidmore (1994), Skidmore (2008)
Sphaeroceroidea: Sphaeroceridae - Lesser Dungflies				
<i>Copromyza equina</i>	Fallén, 1820			Skidmore (2008) cites Roper 1995, Parker 1997
<i>Copromyza nigrina</i>	(Gimmerthal, 1847)			Skidmore (2008)
<i>Crumomyia fimetaria</i>	(Meigen, 1830)			Skidmore (2008)
<i>Crumomyia nigra</i>	(Meigen, 1830)		St Kilda records only	Skidmore (2008)
<i>Crumomyia nitida</i>	(Meigen, 1830)			Skidmore (2008) cites Roper 1995, Parker 1997, 1999

Sphaeroceroidea: Sphaeroceridae - Lesser Dungflies				
<i>Crumomyia pruinosa</i>	(Richards, 1932)	St Kilda records only	Lack (1932), Edwards & Collin (1932), Richards (1932) as <i>C. glacialis</i> var. <i>pruinosa</i> , Skidmore (2008).	
<i>Crumomyia roserii</i>	(Rondani, 1880)		Skidmore (2008) cites Roper 1995	
<i>Herniosina bequaerti</i>	(Villeneuve, 1917)		Skidmore (1994), Skidmore (2008)	
<i>Ischiolepta nitida</i>	(Duda, 1920)		Skidmore (2008) cites Skidmore 1980, Roper 1995	
<i>Leptocera finalis</i>	(Collin, 1956)		Skidmore (1994), Skidmore (2008)	
<i>Leptocera fontinalis</i>	(Fallén, 1826)		Skidmore (1994), Skidmore (2008)	
<i>Leptocera nigra</i>	Olivier, 1813		Skidmore (1994), Skidmore (2008)	
<i>Limosina silvatica</i>	(Meigen, 1830)		Skidmore (2008)	
<i>Lotophila atra</i>	(Meigen, 1830)		Skidmore (1994), Richards (1932) as <i>Copromyza atra</i> , Skidmore (2008) cites Roper 1995, Parker 1997, 1999	
<i>Minilimosina vitripennis</i>	(Zetterstedt, 1847)		Skidmore (1994), Skidmore (2008)	
<i>Opacifrons coxata</i>	(Stenhammar, 1855)		Skidmore (1994), Skidmore (2008)	
<i>Phthitia empirica</i>	(Haliday, 1836)		Skidmore (1994), Skidmore (2008)	
<i>Pseudocollinella humida</i>	(Hutton, 1901)	NBN: <i>Opacifrons humida</i>	Skidmore (1994) as <i>Opacifrons humida</i> , Skidmore (2008).	
<i>Pteremis fenestralis</i>	(Fallén, 1820)		Skidmore (2008)	
<i>Rachispoda fuscipennis</i>	(Haliday, 1833)		Skidmore (2008)	
<i>Rachispoda limosa</i>	(Fallén, 1820)		Skidmore (2008)	
<i>Rachispoda lutosa</i>	Stenhammar, 1855)		Skidmore (1994), Skidmore (2008)	
<i>Spelobia clunipes</i>	(Meigen, 1830)		Skidmore (1994), Skidmore (2008) cites Skidmore 1980	
<i>Spelobia luteilabris</i>	(Rondani, 1880)		Skidmore (2008)	
<i>Spelobia nana</i>	(Rondani, 1880)		Skidmore (2008)	
<i>Spelobia palmata</i>	(Richards, 1927)		Skidmore (2008)	
<i>Spelobia rufilabris</i>	(Stenhammar, 1855)		Skidmore (1994), Skidmore (2008)	
<i>Sphaerocera curvipes</i>	Latreille, 1805		Forrest, Waterston & Watson (1936) as <i>S. subsultans</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)	
<i>Telomerina flavipes</i>	(Meigen, 1830)	St Kilda records only	Skidmore (2008)	
<i>Thoracochaeta brachystoma</i>	(Stenhammar, 1855)		Skidmore (1994), Skidmore (2008)	
<i>Thoracochaeta zosterae</i>	(Haliday, 1833)	<b>Requires review?</b>	Forrest, Waterston & Watson (1936) as <i>Leptocera zosterae</i> , Skidmore (1994), Skidmore (2008) - not checked again Rohacek and Marshall (2000)	



Sphaeroceroidea: Sphaeroceridae - Lesser Dungflies			
<i>Trachyopella minuscula</i>	Collin, 1956		Skidmore (1994), Skidmore (2008)
Ephydroidea: Drosophilidae - Fruit Flies			
<i>Drosophila funebris</i>	(Fabricius, 1787)		Skidmore (2008)
<i>Drosophila phalerata</i>	Meigen, 1830		Skidmore (2008)
<i>Lordiphosa andalusiaca</i>	(Strobl, 1906)	NBN: <i>Drosophila andalusiaca</i>	Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Lordiphosa fenestrarum</i>	(Fallén, 1823)	NBN: <i>Drosophila fenestratum</i>	Skidmore (1994), Skidmore (2008)
<i>Scaptomyza flava</i>	Fallén, 1823		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013, 2019
<i>Scaptomyza graminum</i>	(Fallén, 1823)		Skidmore (1994), Skidmore (2008) cites Roper (1995), Parker 1997, 1999, Bland 1994
<i>Scaptomyza pallida</i>	(Zetterstedt, 1847)		Skidmore (1994), Skidmore (2008)
Ephydroidea: Diastatidae			
<i>Diastata costata</i>	Meigen, 1830		Skidmore (1994), Skidmore (2008)
Ephydroidea: Ephydriidae - Shoreflies			
<i>Hyadina guttata</i>	(Fallén, 1813)		Skidmore (2008)
<i>Hydrellia cochleariae</i>	Haliday, 1839		Skidmore (2008)
<i>Hydrellia maura</i>	Meigen, 1838		Skidmore (1994), Skidmore (2008) cites Parker 1997, 1999 and Roper 1995. Skidmore (2008) - assumes old published records for <i>H. griseola</i> are <i>H. maura</i> . However, <i>H. griseola</i> is a valid taxon and no reason is given for this statement
<i>Hydrellia meigenii</i>	Zatwarnicki, 1988		Skidmore (2008)
<i>Hydrellia ranunculi</i>	Haliday, 1839		Skidmore (2008)
<i>Hydrellia subalbiceps</i>	Collin, 1966		Skidmore (1994), Skidmore (2008) - queries whether specimens from the Uists closer to <i>H. nigricans</i> or whether just a dark form of <i>H. subalbiceps</i> . At the time <i>H. nigricans</i> was not listed by Chandler (2019) and although now included in the UK list there are no records for Scotland on NBN
<i>Lamproscatella sibilans</i>	(Haliday, 1833)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Limnellia quadrata</i>	(Fallén, 1813)		Skidmore (1994)
<i>Limnellia surturi</i>	Andersson, 1971		Skidmore (1994)
<i>Notiphila cinerea</i>	Fallén, 1813		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Notiphila nigricornis</i>	Stenhammar, 1844		Skidmore (2008) cites Roper 1995
<i>Notiphila uglinosa</i>	Haliday, 1839		Skidmore (2008)

Ephydroidea: Ephydridae - Shoreflies			
<i>Parydra coarctata</i>	(Fallén, 1813)		Skidmore (1994), Skidmore (2008)
<i>Parydra pusilla</i>	(Meigen, 1830)		Skidmore (1994), Skidmore (2008)
<i>Parydra quadripunctata</i>	(Meigen, 1830)		Skidmore (2008)
<i>Pelina aenea</i>	(Fallén, 1813)		Skidmore (1994), Skidmore (2008)
<i>Pelina aenescens</i>	(Stenhammar, 1844)		Skidmore (2008)
<i>Pelina nitens</i>	Loew, 1873		Skidmore (1994), Skidmore (2008)
<i>Philygria interstincta</i>	Becker, 1896		Skidmore (1994), Skidmore (2008)
<i>Philygria punctatonervosa</i>	(Fallén, 1813)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013 Mingulay
<i>Scatella paludum</i>	(Meigen, 1830)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Scatella stagnalis</i>	(Fallén, 1813)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Scatella subguttata</i>	(Meigen, 1830)		Skidmore (1994), Skidmore (2008)
<i>Scatophila caviceps</i>	(Stenhammar, 1844)		Skidmore (2008)
Hippoboscoidea: Hippoboscidae - Birdlice-flies			
<i>Melophagus ovinus</i>	(Linnaeus, 1758)		Skidmore (2008); NBN: NTS St Kilda 1986
<i>Ornithomya avicularia</i>	(Linnaeus, 1758)	<b>Requires confirmation</b>	Thompson (1952), Flannan Islands - probably <i>O. fringilla</i> - Grimshaw (1905), Skidmore (2008). Not listed by Waterston (1981)
<i>Ornithomya chloropus</i>	Bergroth, 1901		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2009, 2019
<i>Ornithomya fringillina</i>	Curtis, 1836	<b>Requires confirmation</b>	Thompson (1952)- St Kilda (10/10/1910), Butt Lewis (9/1914) -W. Eagle Clark
Muscoidea: Scathophagidae - Dungflies			
<i>Acanthocnema nigrimana</i>	(Zetterstedt, 1846)		Skidmore (2008)
<i>Ceratinostoma ostiorum</i>	(Haliday in Curtis, 1832)		Whitely, Garland, Hancock (1994), Skidmore (2008); NBN: OHBR 2013-2014
<i>Ernoneura argus</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Gimnomera tarsea</i>	(Fallén, 1819)		Whitely, Garland, Hancock (1994), Bratton (2012), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Norellisoma spinimanum</i>	(Fallén, 1819)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Scathophaga calida</i>	(Haliday in Curtis, 1832)		Forrest, Waterston & Watson (1936) as <i>S. villipes</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: OHBR 2013
<i>Scathophaga furcata</i>	(Say, 1823)		Forrest, Waterston & Watson (1936) as <i>S. squalida</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012-2018
<i>Scathophaga litorea</i>	(Fallén, 1819)		Lack (1932), Whitely, Garland, Hancock (1994), Skidmore (1994), Bratton (2012), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012-2013

Muscoidea: Scathophagidae - Dungflies			
<i>Scathophaga lutaria</i>	(Fabricius, 1794)		Forrest, Waterston & Watson (1936), Skidmore (2008)
<i>Scathophaga stercoraria</i>	(Linnaeus, 1758)		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2019
<i>Scathophaga suilla</i>	(Fabricius, 1794)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2013
<i>Spaziphora hydromyzina</i>	(Fallén, 1819)		Forrest, Waterston & Watson (1936) as <i>Spathiophora hydromyzina</i> , Skidmore (2008)
Muscoidea: Anthomyiidae - Root-maggot Flies			
<i>Anthomyia confusanea</i>	Michelsen in Michelsen & Báez 1985		Skidmore (2008)
<i>Anthomyia liturata</i>	(Robineau-Desvoidy, 1830)		Skidmore (1994), Skidmore (2008) cites Roper (1995)
<i>Botanophila brunneilinea</i>	(Zetterstedt, 1845)		Forrest, Waterston & Watson (1936) as <i>Hylemyia brunneilinea</i> , Skidmore (1994), Skidmore (2008)
<i>Botanophila depressa</i>	(Stein, 1907)		As <i>Pegohylemyia oraria</i> Bland (1994)
<i>Botanophila fugax</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008) cites Roper (1995)
<i>Botanophila seneciella</i>	(Meade, 1892)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Botanophila sericea</i>	(Malloch, 1920)		Skidmore (1994), Skidmore (2008)
<i>Botanophila striolata/discreta agg.</i>	(Fallén, 1824)		As <i>Botanophila striolata</i> , Skidmore (1994), Skidmore (2008) cites Roper (1995)
<i>Chirosia betuleti</i>	(Ringdahl, 1935)	OHBR 2016 first record	NBN: OHBR 2016-2017
<i>Delia albula</i>	(Fallén, 1825)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Delia coarctata</i>	(Fallén, 1825)		Skidmore (2008)
<i>Delia echinata</i>	(Séguy, 1923)		Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Delia florilega</i>	(Zetterstedt, 1845)		Skidmore (2008)
<i>Delia lavata</i>	(Boheman, 1863)	<b>Requires confirmation</b>	Roper & Ackland (2002), Bratton 2010. According to Falk & Pont (2017) it is not possible to distinguish between a small resident breeding population or wind-blown strays from continental Europe
<i>Delia pallipennis</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Delia platura</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Delia quadripila</i>	(Stein, 1916)		Skidmore (2008)
<i>Delia radicum</i>	(Linnaeus, 1758)		Skidmore (2008) cites Roper (1995); NBN: OHBR 2018
<i>Delia setigera</i>	(Stein, 1920)		Skidmore (2008)
<i>Fucellia fucorum</i>	(Fallén, 1819)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper (1995)
<i>Fucellia maritima</i>	(Haliday in Curtis, 1837)		Skidmore (2008); NBN: OHBR 2014

**Muscoidea: Anthomyiidae - Root-maggot Flies**

<i>Fucellia tergina</i>	(Zetterstedt, 1845)		Skidmore (1994), Whitely, Garland, Hancock (1994), Bratton (2012), Skidmore (2008) cites Roper 1995
<i>Hydrophoria linogrisea</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008) cites Roper (1995), Whitely, Garland, Hancock (1994)
<i>Hylemya vagans</i>	(Panzer, [1798])		Skidmore (2008) cites Roper 1995
<i>Hylemya variata</i>	(Fallén, 1823)		Skidmore (1994), Skidmore (2008)
<i>Lasiomma picipes</i>	(Meigen, 1826)		Skidmore (2008) cites Roper 1995
<i>Leucophora cinerea</i>	Robineau-Desvoidy, 1830		Skidmore (2008)
<i>Myopina myopina</i>	(Fallén, 1824)		Skidmore (2008)
<i>Paradelia brunneonigra</i>	(Schnabl in Schnabl & Dziedzicki, 1911)		Skidmore (2008) cites Bratton 1987
<i>Paradelia intersecta</i>	Meigen, 1826)		Skidmore (1994), Skidmore (2008)
<i>Paregle audacula</i>	(Harris, [1780])		Skidmore (1994), Skidmore (2008)
<i>Pegomya betae</i>	(Curtis, 1847)		Skidmore (2008) cites Skidmore 1989
<i>Pegomya bicolor</i>	(Wiedemann, 1817)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper (1995); NBN: OHBR 2018
<i>Pegomya fulgens</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008)
<i>Pegomya haemorrhoum</i>	(Zetterstedt, [1838])		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Pegomya hyoscyami</i>	(Panzer, [1809])		Bland (1994), Skidmore (2008) cites Bratton 1987
<i>Pegomya solennis</i>	(Meigen, 1826)		Skidmore (2008) cites Skidmore 1989, Roper 1995: NBN: OHBR 2017
<i>Pegoplata aestiva</i>	(Meigen, 1826)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Pegoplata infirma</i>	(Meigen, 1826)		Skidmore (1994), Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Phorbia fumigata</i>	(Meigen, 1826)		Skidmore (2008)
<i>Subhylemyia longula</i>	(Fallén, 1824)		Skidmore (1994), Skidmore (2008)
<i>Zaphne ambigua</i>	(Fallén, 1823)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Zaphne caudata</i>	(Zetterstedt, 1855)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Dinnin 1999

**Muscoidea: Fanniidae - Lesser Houseflies**

<i>Fannia canicularis</i>	(Linnaeus, 1761)		Skidmore (1994), Skidmore (2008); NBN: OHBR 2015
<i>Fannia fuscula</i>	(Fallén, 1825)		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008)
<i>Fannia genualis</i>	(Stein, 1895)		Skidmore (1994), Skidmore (2008)
<i>Fannia incisurata</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Fannia lepida</i>	(Zetterstedt, 1845)		Skidmore (1994), Skidmore (2008)

<i>Fannia mollissima</i>	(Haliday in Westwood, 1840)	Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Fannia parva</i>	(Stein, 1895)	Waterston (1981) new record South Uist, Skidmore (2008)
<i>Fannia postica</i>	(Stein, 1895)	Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Fannia rondanii</i>	(Strobl, 1893)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Fannia serena</i>	(Fallén, 1825)	Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<b>Muscoidea: Muscidae - Houseflies</b>		
<i>Azelia cilipes</i>	(Haliday, 1838)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Azelia gibbera</i>	(Meigen, 1826)	Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Azelia zetterstedti</i>	Rondani, 1866	Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Coenosia albicornis</i>	Meigen, 1826	Skidmore 2008
<i>Coenosia campestris</i>	(Robineau-Desvoidy, 1830)	Skidmore 2008
<i>Coenosia femoralis</i>	(Robineau-Desvoidy, 1830)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Coenosia lacteipennis</i>	(Zetterstedt, 1845)	Forrest, Waterston & Watson (1936) a <i>Dexiopsis lactipennis</i> , Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Coenosia means</i>	Meigen, 1826	Forrest, Waterston & Watson (1936), Skidmore 2008
<i>Coenosia mollicula</i>	(Fallén, 1825)	Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Coenosia pedella</i>	(Fallén, 1825)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper
<i>Coenosia pumila</i>	(Fallén, 1825)	Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Coenosia rufipalpis</i>	Meigen, 1826	Skidmore 2008
<i>Coenosia testacea</i>	(Robineau-Desvoidy, 1830)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Coenosia tigrina</i>	(Fabricius, 1775)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2012-2018
<i>Drymeia hamata</i>	(Fallén, 1823)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: OHBR 2017, 2018
<i>Eudasyphora cyanella</i>	(Meigen, 1826)	Skidmore 2008
<i>Graphomya maculata</i>	(Scopoli, 1763)	Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008)
<i>Graphomya minor</i>	Robineau-Desvoidy, 1830	Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Hebecnema umbratica</i>	(Meigen, 1826)	Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Hebecnema vespertina</i>	(Fallén, 1823)	Skidmore (2008) cites Roper 1995; NBN: OHBR 2015
<i>Helina arctata</i>	Collin, 1953	Whitely, Garland, Hancock (1994), Skidmore (2008)

Muscoidea: Muscidae - Houseflies			
<i>Helina calceata</i>	(Rondani, 1866)		Forrest, Waterston & Watson (1936) as <i>Mydaea calceata</i> , Skidmore (2008)
<i>Helina cinerella</i>	(van der Wulp, 1867)		Skidmore (2008) cites Roper 1995
<i>Helina confinis</i>	(Fallén, 1825)		Skidmore (2008)
<i>Helina cothurnata</i>	(Rondani, 1866)		Skidmore (2008)
<i>Helina evecta</i>	(Harris, 1780)		Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2015
<i>Helina impuncta</i>	(Fallén, 1825)		Forrest, Waterston & Watson (1936) as <i>Mydaea impuncta</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Helina maculipennis</i>	(Zetterstedt, 1845)		Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2015
<i>Helina obscurata</i>	(Meigen, 1826)		Skidmore (2008)
<i>Helina protuberans</i>	(Zetterstedt, 1845)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995
<i>Helina pubiseta</i>	(Zetterstedt, 1845)		Skidmore (2008)
<i>Helina quadrum</i>	(Fabricius, 1805)		Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Helina reversio</i>	(Harris, [1780])		Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2015
<i>Helina setiventris</i>	Ringdahl, 1924		Skidmore (2008) cites Roper 1995
<i>Helina subvittata</i>	(Séguy, 1923)		Skidmore (2008)
<i>Helina trivittata</i>	(Zetterstedt, 1860)		Skidmore (2008)
<i>Hydrotaea armipes</i>	(Fallén, 1825)		Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Hydrotaea cinerea</i>	Robineau-Desvoidy, 1830		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Hydrotaea dentipes</i>	(Fabricius, 1805)		Forrest, Waterston & Watson (1936), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2018
<i>Hydrotaea irritans</i>	(Fallén, 1823)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2015, 2018
<i>Hydrotaea similis</i>	Meade, 1887	OHBR 2013 first record	NBN: OHBR 2013
<i>Limnophora maculosa</i>	(Meigen, 1826)		Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Limnophora olympiae</i>	Lyneborg, 1965		Skidmore (2008)
<i>Limnophora triangula</i>	(Fallén, 1825)		Skidmore (2008)
<i>Limnospila albifrons</i>	(Zetterstedt, 1849)		Skidmore (1994), Skidmore (2008)
<i>Lispe tentaculata</i>	(Zetterstedt, 1849)		Skidmore (2008)
<i>Morellia hortorum</i>	(Fallén, 1817)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2015, 2018

Muscoidea: Muscidae - Houseflies			
<i>Morellia simplex</i>	(Loew, 1857)	Requires review	NBN: BRC 1910, 1936, 1966 - not listed by Waterston (1981) or Skidmore (2008)
<i>Musca domestica</i>	Linnaeus, 1758	Requires review	Grimshaw 1905 - Flannan Islands W. Eagle Clarke, Skidmore (2008). A single record from the Flannan Islands
<i>Mydaea anicula</i>	(Zetterstedt, 1860)	OHBR 2015 first record	NBN: OHBR 2015
<i>Mydaea urbana</i>	(Meigen, 1826)		Skidmore (2008)
<i>Myospila mediatubunda</i>	(Fabricius, 1781)		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2017
<i>Neomyia cornicina</i>	(Fabricius, 1781)		Forrest, Waterston & Watson (1936) as <i>Cryptolucilia caesarion</i> , Skidmore (1994), Skidmore (2008)
<i>Phaonia angelicae</i>	(Scopoli, 1763)		Forrest, Waterston & Watson (1936) as <i>P. basalis</i> , Skidmore (2008); NBN: OHBR 2018
<i>Phaonia consobrina</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Phaonia errans</i>	(Meigen, 1826)		Skidmore (2008)
<i>Phaonia erronea</i>	(Schnabl, 1887)	OHBR 2015 first record	NBN: OHBR 2015
<i>Phaonia incana</i>	(Wiedemann, 1817)		Forrest, Waterston & Watson (1936), MacGowan (1994) Flannan Islands, Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper; NBN: OHBR 2015-2018
<i>Phaonia perdita</i>	(Meigen, 1830)		Skidmore (2008) cites Roper 1995; NBN: OHBR 2019
<i>Pseudocoenosia abnormis</i>	Stein, 1916		Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Schoenomyza litorella</i>	(Fallén, 1823)		Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008)
<i>Spilogona aerea</i>	(Fallén, 1825)		Skidmore (2008)
<i>Spilogona contractifrons</i>	(Zetterstedt, 1838)	OHBR 2015 first record	NBN: OHBR 2015
<i>Spilogona denigrata</i>	(Meigen, 1826)		Forrest, Waterston & Watson (1936) as <i>Limnophora denigrata</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Spilogona falleni</i>	Pont, 1984		Skidmore (2008)
<i>Spilogona litorea</i>	(Fallén, 1823)		Forrest, Waterston & Watson (1936) as <i>Limnophora litorea</i> , Waterston (1981), Whitely, Garland, Hancock (1994)
<i>Spilogona pacifica</i>	(Meigen, 1826)		Whitely, Garland, Hancock (1994) as <i>S. varda</i> , Skidmore (2008)
<i>Spilogona solitaria</i>	(Collin, 1921)		Skidmore (2008)
<i>Spilogona surda</i>	(Zetterstedt, 1845)		Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Spilogona trianguligera</i>	(Zetterstedt, 1838)		Skidmore (2008)
<i>Thricops cunctans</i>	(Meigen, 1826)		Skidmore (2008)
<i>Thricops foveolatus</i>	(Zetterstedt, 1845)		Skidmore (2008)



Muscoidea: Muscidae - Houseflies				
<i>Thricops longipes</i>	(Zetterstedt, 1845)	St Kilda records only		Skidmore (2008)
<i>Thricops nigritellus</i>	(Zetterstedt, 1838)			Skidmore (2008)
<i>Thricops rostratus</i>	(Meade, 1882)			Skidmore (2008)
<i>Thricops semicinereus</i>	(Wiedemann, 1817)			Skidmore (2008)
Oestroidea: Calliphoridae - Blowflies				
<i>Bellardia pandia</i>	(Walker, 1849)	OHBR 2013 first record		NBN: OHBR 2013
<i>Bellardia pubicornis</i>	(Zetterstedt, 1838)			Skidmore (2008)
<i>Bellardia viarum</i>	(Robineau-Desvoidy, 1830)	OHBR 2017 first record		NBN: OHBR 2017
<i>Bellardia vulgaris</i>	(Robineau-Desvoidy, 1830)			Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013. According to Skidmore (2008) old published records of <i>Onesia sepulchralis</i> (Linnaeus) from the Uists presumably refer to <i>B. vulgaris</i> but some records may relate to <i>B. viarum</i>
<i>Calliphora loewii</i>	Enderlein, 1903			Whitely, Garland, Hancock (1994), Skidmore (2008)
<i>Calliphora subalpina</i>	(Ringdahl, 1931)	OHBR 2014 first record		NBN: OHBR 2014
<i>Calliphora uralensis</i>	Villeneuve, 1922			Skidmore (1994), Bratton (2012), Skidmore (2008), MacDonald (2014), MacDonald (2016); NBN: OHBR 2010-2018
<i>Calliphora vicina</i>	Robineau-Desvoidy, 1830			Forrest, Waterston & Watson (1936) as <i>C. erythrocephala</i> , Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2019
<i>Calliphora vomitoria</i>	(Linnaeus, 1758)			Skidmore (1994), Skidmore (2008); NBN: OHBR 2013-2018
<i>Cynomya mortuorum</i>	(Linnaeus, 1761)			Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2019
<i>Lucilia caesar</i>	(Linnaeus, 1758)			Skidmore (1994), Skidmore (2008)
<i>Lucilia illustris</i>	(Meigen, 1826)			Skidmore (1994), Skidmore (2008) cites Roper 1995
<i>Lucilia sericata</i>	(Meigen, 1826)			Forrest, Waterston & Watson (1936), Whitely, Garland, Hancock (1994), Skidmore (1994), Skidmore (2008); NBN: OHBR 2013-2018
<i>Lucilia silvarum</i>	(Meigen, 1826)	OHBR 2013 first record		NBN: OHBR 2013
<i>Melinda gentilis</i>	Robineau-Desvoidy, 1830			Skidmore (2008) as <i>Onesia caerulea</i>
<i>Melinda viridicyanea</i>	(Robineau-Desvoidy, 1830)			Skidmore (2008)
<i>Protophormia terraenovae</i>	(Robineau-Desvoidy, 1830)			Skidmore (2008); NBN: OHBR 2013-2017

<b>Oestroidea: Rhinophoridae</b>			
<i>Melanophora roralis</i>	(Linnaeus, 1758)		Skidmore (2008)
<b>Oestroidea: Sarcophagidae - Fleshflies</b>			
<i>Sarcophaga melanura</i>	Meigen, 1826		Skidmore (2008)
<i>Sarcophaga nigriventris</i>	Meigen, 1826		Skidmore (2008) cites Roper 1995
<b>Oestroidea: Tachinidae - Parasite-flies</b>			
<i>Campylocheta inepta</i>	(Meigen, 1824)		Skidmore (2008) cites Skidmore 1989
<i>Ceromya bicolor</i>	(Meigen, 1824)		Skidmore (1994), Skidmore (2008)
<i>Eurithia anthophila</i>	(Robineau-Desvoidy, 1830)		Skidmore (2008) cites Parker 2007; NBN: BRC 2002
<i>Gymnocheta viridis</i>	(Fallén, 1810)		Whitely, Garland, Hancock (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013
<i>Linnaemya vulpina</i>	(Fallén, 1810)		Skidmore (1994), Skidmore (2008) cites Parker 1999, 2004
<i>Lydina aenea</i>	(Meigen, 1824)		Skidmore (2008) cites Roper 1995
<i>Medina collaris</i>	(Fallén, 1820)		Skidmore (1994), Skidmore (2008) cites Roper 1995; NBN: OHBR 2013-2018
<i>Pales pavidata</i>	(Meigen, 1824)		Skidmore (2008)
<i>Pelatachina tibialis</i>	(Fallén, 1810)	OHBR 2015 first record	NBN: OHBR 2015
<i>Phryxe</i>	Robineau-Desvoidy, 1830	OHBR 2013 first record	NBN: OHBR 2013
<i>Platymya fimbriata</i>	(Meigen, 1824)	BRC 2009 - First record	NBN: BRC 2009
<i>Prosenia siberita</i>	(Fabricius, 1775)		Forrest, Waterston & Watson (1936) as <i>P. luculliana</i> , Whitely, Garland, Hancock (1994), Skidmore (2008) cites Parker 2002, 2004; NBN: BRC 2009,
<i>Siphona geniculata</i>	(De Geer, 1776)		Forrest, Waterston & Watson (1936) as <i>Bucentes geniculata</i> , Skidmore (2008) cites Roper 1995
<i>Siphona maculata</i>	Staeger in Zetterstedt, 1849		Skidmore (2008) cites Roper 1995
<i>Tachina grossa</i>	(Linnaeus, 1758)	OHBR 2019 first record	NBN: OHBR 2019
<i>Winthemia quadripustulata</i>	(Fabricius, 1794)	<b>Not confirmed</b>	Waterston 1981), recorded as <i>Winthemia (Exorista) apicalis</i> (Dale 1892) Vallay, North Uist
<b>Oestroidea: Oestridae - Warble-flies, Bots</b>			
<i>Gasterophilus intestinalis</i>	(De Geer, 1776)		Skidmore (2008)
<i>Hypoderma lineatum</i>	(De Villers, 1789)		Forrest, Waterston & Watson (1936) – “Cattle infested by warbles were frequently seen on the machair” (Barra), Skidmore (2008)